

Research Paper



Bioactive compounds of prosopis africana oil (african mesquite) using gas chromatography and mass spectrometry (gc-ms) technique

Alagbe, John Olujimi*

*Department of Animal Science, University of Abuja, Nigeria.

Article Info

Article History:

Received: 26 December 2023

Revised: 03 March 2023

Accepted: 11 March 2023

Published: 27 April 2023

Keywords:

Phytochemicals

Prosopis Africana

Pharmacological Properties

Gas Chromatography

Antiarrhythmic

Antioxidant



ABSTRACT

Phytochemicals (bioactive compounds) are generally regarded as chemicals of plant origin used for plants for growth, defense against competitors and pathogens. They also have a wide range of pharmacological activities including – antiarrhythmic, antioxidant, antimalarial, vasodilatory, antimicrobial, analgesics, hepato-protective, anti-inflammatory, hypo-cholesterolemic, anti-androgenic, antiviral, antifungal and immune-modulatory. This study was carried out to investigate the bioactive compounds of Prosopis africana oil (African mesquite) using gas chromatography – mass spectrometry technique. A total of 73 bioactive compounds (77.16 %) were identified based on their peak areas. The major compounds identified in the oil sample were; prosogerin A (29.90 %), caryophyllene (12.33 %), 2, 4-bis (1-phenylethyl) phenol (5.80 %), gallic acid (5.22 %), β -cyclocitral (3.11 %), β -sitosterol (2.50 %), α -terpinene (2.09 %) and ellagic acid (2.04 %). The other compounds were less than 1 %, however, they exert various pharmacological properties. It was concluded that Prosopis africana oil is loaded with phytochemicals and also has several health promoting properties and it can also be used as a remedy for the treatment of several ailments in human being and animals.

Corresponding Author:

Alagbe, John Olujimi

Department of Animal Science, University of Abuja, Nigeria.

Email: dralagbe@outlook.com

Copyright © 2023 The Author(s). This is an open access article distributed under the Creative Commons Attribution License, (<http://creativecommons.org/licenses/by/4.0/>) which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

1. INTRODUCTION

Prosopis africana (African mesquite) is an evergreen multipurpose leguminous plant belonging to the family Fabaceae. It has about 45 species and it is widely distributed in most part of Africa, Asia and America [1], [2].

The seeds of *Prosopis africana* are rich in phyto-constituents such as: alkaloids, tannins, flavonoids, terpenoids, steroids, saponins, glycosides and phenols with significant therapeutic properties [3], [4], [5], [6].

The seeds has antioxidant, antifungal, antimicrobial, antiparasitic, analgesics, antipyretic, anti-proliferative, antifungal, anti-inflammatory, hepato-protective, antiprotozoal, anti-androgenic and cytotoxic properties [7], and it also contains important minerals such as calcium, potassium, zinc, manganese, magnesium, copper and sodium [8], [9], [5], [10], [11]. These minerals are vital for the activation of key enzymes that provide a greater protection from diseases [12], [13], [14], [15], [16], [17], [18], [19].

Moreover, *Prosopis africana* seeds contains amino acid alanine, proline, valine and isoleucine which are the main structural protein found in connective tissues of animals [20], [21], [22]. The medicinal properties of plants depends on its composition of phyto-constituents or bioactive compounds [23], [24].

The stem bark and root decoction of *Prosopis* are used traditionally for the treatment of rheumatism, cough, malaria, bronchitis, skin diseases, gastro intestinal and sexually transmitted infections [25]. Fermented seeds of the plant can be used in the preparation of soup because of its sweet smell [26], [27], [28], [29], [30].

Preclinical investigations has shown the pharmacological responses of *Prosopis africana* oil to inhibit the activities of some pathogenic organisms such as; *Salmonella* spp, *Candida albicans*, *Escherichia coli*, *Bacillus subtilis*, *Bacillus pumilus*, *Bacillus licheniformis* and *Bacillus megaterium* [31]. Aqueous and ethanolic extracts from the leaves and stem bark of *Prosopis africana* can be used for the treatment of diarrhea and coccidiosis in poultry [32].

Identifying the bioactive compounds in a medicinal plant could lead to the production of novel drugs [18], [33]. One key methods in identifying unknown compounds, quantification and characterization is the use of gas chromatography-mass spectrometry (GC-MS) technique.

Investigating the bioactive compounds in a sample will give a clue on its efficacy and safety. Hence, in view of the immense medicinal properties in *Prosopis africana* seed. This study was designed to evaluate the phytochemicals or bioactive compounds in *Prosopis africana* oil using gas chromatography-mass spectrometry (GC-MS) technique.

2. METHODOLOGY

2.1 Experimental Site

This study was carried out at the Department of Animal Science Laboratory, University of Abuja main campus along Airport road, Gwagwalada, Abuja, Nigeria located between latitudes 8°57'N and 8°55'N and longitude 7°05'E and 7°06'E [34].

2.2 Collection, Identification Processing of *Prosopis Africana* Oil

Prosopis africana seed were purchased from a local market in Gwagwalada, Nigeria. It was identified and authenticated at the Herbarium of Department of Biological Sciences, University of Abuja, Nigeria with a voucher number ABJ/09A/20. It was shade dried for 13 days and grinded into powder using a laboratory blender (Panasonic, Model EDH-023) and stored in a labeled polythene bag for further test

2.3 Extraction of *Prosopis Africana* Oil Using Steam Distillation Method

Prosopis africana oil was extracted using steam-distillation technique, the entire extraction procedure requires: digital weighing scale, round bottom flask, distilled water, glass yarn heating mantle, measuring cylinder, separatory funnel. 50 grams of grinded *Prosopis* seed was weighed into a round

bottom flask, mixed with 250 mL of distilled water. The mixture was placed in a glass yarn heating mantle, adjusted to a temperature of 80 °C before setting up the condenser above the round bottom flask.

The mixture is boiled vigorously for 15 minutes and the distillate was collected in a beaker until no more droplets of oil come over followed by the addition of 5 grams of sodium chloride and stirred vigorously. Thereafter, the distillate was transferred into the separatory funnel and evaporated in a steam bath to obtain *Prosopis africana* oil.

2.4 Bioactive Compounds of PRAO Using GC-MS Technique

Bioactive compounds of PRAO were identified using LABTRON GC-MS (GC-MS-879). It is a high precision gas chromatography mass spectrometer with pre-filter mass analyzer and electron multiplier ensuring high sensitivity.

Gas chromatography (Model-GC-87) has the following technical specifications; flow rate (1-15 ml / min), maximum flow rate (10 ml/min), inlet temperature (450° C), pressure range (0 – 100 psi), heating rate (up to 120° C/min), room temperature (4° C-450° C), pressure control mode (electronic pressure control), split mode (split / split less), split ratio (1000: 1) and temperature programming (7 stages / 8 platforms) while the mass spectrometry (Model-MS-89): mass range (1.5-1000 amu), ion-source temperature (100 ° C-350° C), GC-MS interface temperature (450° C), scan rate (up to 1000 amu/sec), stability (\pm 0.10 amu/ 48 hours), filament emission current (0 - 350° C μ A), EI source ionization energy (5 eV-250 eV), sensitivity (S/N is \geq 30:1) and vacuum (Turbo molecular pump: 67 L/s).

Volatile compounds were identified with standard compounds in National Institute of Standard and Technology mass spectral library using Chemstation library database. Percentage area were based on the ratio between the peak area of each compound and the sum of the peak areas of all compounds.

Table 1. Bioactive Compounds of *Prosopis Africana* Oil as Detected by GC-MS

Peak No	RT (min)	Area (%)	Name of Compound
1	2.77	29.90	Prosogerin A (6 Methoxy-7-hydroxyl dioxyflavone)
2	3.93	5.80	2,4-bis (1-phenylethyl) phenol
3	4.03	0.63	β -phenethylamine
4	4.51	0.14	2,4,6-tris(1-phenylethyl) phenol
5	4.92	0.16	α -pinene
7	5.22	2.09	α -terpinene
8	5.91	0.46	β -pinene
9	6.07	0.12	β -myrcene
10	6.34	0.74	α -phellandrene
11	6.88	0.11	α -terpinolene
12	7.22	0.35	γ -terpinene
13	8.09	0.03	1-terpineol
14	8.44	0.07	4-terpineol
15	9.57	0.18	Humulene
16	9.88	12.33	Caryophyllene
17	10.12	1.85	Copaene
18	11.55	0.21	Cis-linaloxide
19	11.83	0.30	α -Selinene
20	11.90	0.06	γ -Elemene
21	12.10	0.01	α -Gurjunene
22	12.33	0.42	β -Elemene
23	12.68	3.11	β -Cyclocitral
24	12.75	0.05	3-hexenyl-2-methylbutanoate
25	12.92	0.02	Exo-methyl-camphenilol
26	13.10	0.06	Caryophyllene oxide

27	13.47	0.03	Benzene -1-methoxy-2-methyl
28	13.55	0.18	Napthalene, 1,2 hydro-1,1, 6 -trimethyl
29	13.80	1.55	Cycloheptasiloxane, tetradecamethyl
30	13.87	1.41	Nerolidyl acetate
31	13.91	0.001	1-Cyclohexene-1- butanal, alpha, 2, 6, 6- tetramethyl
32	14.00	0.11	Cyclooctasiloxane, hexadecamethyl
33	14.31	0.60	2,2, 6-Trimethyl-2H, 5Hpyrano(3,2-c) quinolin-5one
34	14.49	0.27	Dibutyl phthalate
35	14.52	0.70	Acetic acid, dodecyl ester
36	14.70	0.64	1,1, 5-Trimethyl-1,2-dihydronaphthalene
37	14.83	0.02	Alloaromedendrene oxide
38	14.92	0.03	Butylated Hydroxytoluene
39	14.95	0.01	Octacosane
40	16.71	0.16	Kaur-16-ene
41	16.37	0.10	1,5,8-p-Menthatriene
42	16.55	0.20	Spicigerine
43	18.28	0.07	Prosophylline
44	18.84	2.04	Ellagic acid
45	18.92	0.55	Apigenin-6-glucoside
46	19.20	1.80	Quercetin-3-rhamnoside
47	19.44	0.01	Isoprosopilosine
48	19.75	1.50	Prosopilosidine
49	19.87	0.02	Trimethyl-tetrahydronaphthalene
50	21.21	0.21	Spathulenol
51	21.40	0.01	1,3-benzodioxole,5-(2,2-dimethyl)
52	21.61	0.22	2-nonen-4-one
53	22.02	0.11	Bicyclo [2.2.1] heptan-3-one
54	25.04	2.50	β -sitosterol
55	25.80	0.32	Cis-vaccenic acid
56	26.11	0.16	3-chloro-N-isochroman-1-methyl propionamide
57	26.70	0.03	3,4 - dichlorobenzonitrile
58	26.84	0.07	Mannitol, 1,4-di-O-methyl tetraacetate
59	27.02	0.14	β -1-rhamnofuranoside thio-octyl
60	27.18	0.03	Guaia-1(10), 11-diene
61	27.35	0.12	3,4 - dimethyl phenyl heptyl ether
62	27.80	0.09	1,1,5 -trimethyl -1,2 -dihydronaphthalene
63	27.93	0.03	1-cyclohexane-1-propanal
64	28.06	0.04	1-Trimethylsilylpent-1-en-4-yne
65	28.93	0.02	β -ocimene
66	28.98	0.05	Trans- β -ocimene
67	33.40	0.18	1-Methyl-6-methylenebicycloheptane
68	33.73	0.04	1,6,10,14 -Hexadecatetraen-3-ol
69	33.81	0.01	Alloaromadendrene oxide (1)
70	33.92	0.40	1-cyclohexane-1-propanal
71	36.27	0.18	1-Formyl -2,2,6 -trimethyl cyclohexane
72	36.84	5.22	Gallic acid
73	36.93	0.06	6-Isopropenyl-4,8-dimethyl nephthalen-2-ol
74	37.10	0.01	1,6,10-Dodecatrien-3-trimethyl -ol
Total		77.16	

RT: Reaction time (min)

3. RESULTS AND DISCUSSION

3.1 Bioactive Compounds of Prosopis Africana Oil

Bioactive compounds of Prosopis africana oil is presented in Table 1 a total of 73 bioactive compounds (77.16 %) were identified based on their peak areas. The major compounds identified in the oil sample were; prosogerin A (29.90 %), caryophyllene (12.33 %), 2, 4-bis (1-phenylethyl) phenol (5.80 %), gallic acid (5.22 %), β -cyclocitral (3.11 %), β -sitosterol (2.50 %), α -terpinene (2.09 %) and ellagic acid (2.04 %). Other minor compounds includes; β -phenethylamine (0.63 %), 2, 4, 6-tris (1-phenylethyl) phenol (0.14 %), α -pinene (0.16 %), β -myrcene (0.12 %), α -phellandrene (0.74 %), α -terpinolene (0.11 %), γ -terpinene (0.35 %), 1-terpineol (0.03 %), 4-terpineol (0.07 %), humulene (0.18 %), copaene (0.85 %), cis-linaloxide (0.21 %), α -selinene (0.30 %), γ -elemene (0.06 %), α -gurjunene (0.01 %), β -elemene (0.42 %), 3-hexenyl-2-methylbutanoate (0.05 %), exo-methyl-camphenilol (0.02 %), benzene -1-methoxy-2-methyl (0.03 %), naphthalene, 1,2 hydro-1,1, 6 -trimethyl (0.18 %), cycloheptasiloxane, tetradecamethyl (1.55 %), nerolidyl acetate (1.41 %), 1-cyclohexene-1- butanal, alpha, 2, 6, 6- tetramethyl (0.001 %), cyclooctasiloxane, hexadecamethyl (0.11 %), 2,2, 6-trimethyl-2H, 5H pyrano (3,2-c) quinolin-5-one (0.60 %), dibutyl phthalate (0.27 %), acetic acid, dodecyl ester (0.70 %), 1,1, 5-trimethyl-1,2-dihydronaphthalene (0.64 %), alloaromadendrene oxide (0.02 %), butylated hydroxytoluene (0.03 %), octacosane (0.01 %), kaur-16-ene (0.16 %), 1,5,8-p-menthatriene (0.10 %), spicigerine (0.20 %), prosophylline (0.07 %), apigenin-6-glucoside (0.55 %), quercetin-3-rhamnoside (1.80 %), isosprosopilosine (0.01 %), prosopilosidine (1.50 %), trimethyl-tetrahydronaphthalene (0.02 %), spathulenol (0.21 %), 1,3-benzodioxole,5-(2,2-dimethyl) (0.01 %), 2-nonen-4-one (0.22 %), bicyclo [2.2.1] heptan-3-one (0.11 %), cis-vaccenic acid (0.32 %), 3-chloro-N-isochroman-1-methyl propionamide (0.16 %), 3,4 - dichlorobenzonitrile (0.03 %), mannitol, 1,4-methyl tetraacetate (0.07 %), β -1-rhamnofuran oside thio-octyl (0.14 %), guaia-1(10), 11-diene (0.03 %), 3,4 - dimethyl phenyl heptyl ether (0.12 %), 1,1,5 -trimethyl -1,2 -dihydronaphthalene (0.09 %), 1-cyclohexane-1-propanal (0.03 %), 1-trimethylsilylpent-1-en-4-yne (0.04 %), β -ocimene (0.02 %), trans- β -ocimene (0.05), -methyl-6-methylenebicycloheptane (0.18 %), 1,6,10,14 -hexadecatetraen-3-ol (0.04 %), alloaromadendrene oxide (1) (0.01 %), 1-cyclohexane-1-propanal (0.40 %), 1-formyl -2,2,6 -trimethyl cyclohexane (0.18 %), 6-isopropenyl-4,8-dimethyl naphthalen-2-ol (0.06 %) and 1,6,10-dodecatrien-3-trimethyl -ol (0.01 %). However all these compounds have significant pharmacological effects [35], [36].

Prosogerin A has been reported to possess antimicrobial and anti-inflammatory properties [5]. Caryophyllene occurs naturally in essential oils of several plants including; Piper nigrum, Lantana camara leaf oil and Capaifera spp and has been reported to function as antioxidant, thus scavenging free radicals (Adane, 2020; Pagare et al., 2015). 2,4-bis (1-phenylethyl) phenol, α -terpinene, 2,4,6-tris(1-phenylethyl) phenol, β -myrcene, α -terpinolene and α -phellandrene had been reported by researchers to have anti-inflammatory and hypocholesterolemic properties [37], [38]. Humulene, β -sitosterol, α -gurjunene, α -selinene and exo-methyl-camphenilol possess hepato-protective, anti-proliferative, antifungal, antipyretic and antihelminthic properties [11], [12], [5], [10], [11], [38]. Naphthalene, 1,2 hydro-1,1, 6 - trimethyl and benzene -1-methoxy-2-methyl are capable of performing nematicide and anti-androgenic activities [40].

The other bioactive compounds in less concentration contained in Prosopis africana oil also have significant therapeutic properties and are used in ayurvedic remedies for a variety of ailments [36]. However, the result obtained in this experiment is in agreement with the findings of [41], [38], [42], [43], but contrary to the reports of [5]. This differences are influenced by factors like species, climate, harvest stage, location, storage condition and method of extraction of the oil [44], [45], [46], [47], [48], [49], [50] [24].

4. CONCLUSION

It can be concluded that *Prosopis africana* oil possess several health promoting properties (antimicrobial, antifungal, antiviral, anti-fibrotic, immune-modulatory, cytotoxic, antipyretic, antitumor, antihelminthic and hepato-protective) due to the presence of bioactive compounds. These compounds can reduce the rising concern on environmental pollution, antibiotic resistance and food safety. It could also enhance livestock performance and production of novel drugs.

Acknowledgments

The authors have no specific acknowledgments to make for this research.

Funding Information

This research received no specific grant from any funding agency in the public, commercial, or not-for-profit sectors.

Author Contributions Statement

Name of Author	C	M	So	Va	Fo	I	R	D	O	E	Vi	Su	P	Fu
Alagbe, John Olujimi	✓	✓	✓	✓		✓		✓	✓	✓	✓			

C : Conceptualization

M : Methodology

So : Software

Va : Validation

Fo : Formal analysis

I : Investigation

R : Resources

D : Data Curation

O : Writing - Original Draft

E : Writing - Review & Editing

Vi : Visualization

Su : Supervision

P : Project administration

Fu : Funding acquisition

Conflict of Interest Statement

The authors declare that there are no conflicts of interest regarding the publication of this paper.

Informed Consent

All participants were informed about the purpose of the study, and their voluntary consent was obtained prior to data collection.

Ethical Approval

The study was conducted in compliance with the ethical principles outlined in the Declaration of Helsinki and approved by the relevant institutional authorities.

Data Availability

The data that support the findings of this study are available from the corresponding author upon reasonable request.

REFERENCES

- [1] M. U. Adikwu, S. Ezeabasili, and C. O. Esimone, 'Evaluation of the physico- chemical properties of a new polysaccharide gum from *Prosopis africana*', *Bollettino Chimico Farmaceutico*, vol. 140, no. 1, pp. 40-45, 2001. doi.org/10.4314/dai.v12i1.15571
- [2] A. O. Ugwoke, S. Oluka, and P. C. Eze, 'Physicomechanical properties of iron tree seed (*Prosopis africana*) from South Eastern Nigeria', *Journal of Engineering and Technology*, vol. 24, no. 1, pp. 13-18, 2019.
- [3] J. O. Alagbe, 'Prosopis africana stem bark as an alternative to antibiotic feed additives in broiler chick's diets: Performance and Carcass characteristics', *Journal of Multidimensional Research and Reviews*, vol. 2, no. 1, pp. 64-77, 2021.


- [4] J. O. Alagbe, M. O. Adedeji, Z. Habiba, G. Nwosu, and W. Dabara Comfort, 'Physico-chemical properties of *Indigofera zollingeriana* seed oil', *Asian Journal of Advances in Medical Science*, vol. 3, no. 4, pp. 306-308, 2021.
- [5] O. C. P. Agubosi, R. Oluwafemi, and J. O. Alagbe, 'Preliminary Study on GC-MS Analysis of *Prosopis africana* Seed (African Mesquite) oil', *Journal of Ethics and Diversity in International Communication*, vol. 1, no. 4, pp. 18-21, 2021.
- [6] K. S. Olorumaiye, L. E. Apeh, H. Madandola, and M. O. Oguntoye, 'Proximate and phytochemical composition of African mahogany seed and African mesquite pod', *Journal of Applied Sciences and Environmental Management*, vol. 23, no. 2, pp. 249-252, 2019. doi.org/10.4314/jasem.v23i2.8
- [7] S. Harzallah, 'Flavonoids diversification in organs of two *Prosopis fraxta* (Bank and Sol.) eig. (Leguminosea, Mimosoidae) populations occurring in Northeast and the Southeast of Tunisia', *Journal of Applied Research*, vol. 1, pp. 130-136, 2005.
- [8] L. R. Ferguson and P. Harris, 'Antioxidant and antigenotoxic effects of plant cell wall hydroxycinnamic acids in cultured HT-29 cells', *Molecular Nutrition and Food Research*, vol. 49, pp. 585-593, 2005. doi.org/10.1002/mnfr.200500014
- [9] S. Pagare, M. Bhatia, and N. Tripathi, 'Secondary metabolites of plants and their role: Overview', *Overview. Current Trends in Biotechnology Pharmacology*, vol. 9, pp. 294-305, 2015.
- [10] R. A. Oluwafemi, O. C. Agubosi, and J. O. Alagbe, 'Proximate, minerals, vitamins and amino acid composition of *Prosopis africana* (African mesquite) seed oil', *Asian Journal of Advances in Research*, vol. 11, no. 1, pp. 21-27, 2021.
- [11] O. C. P. Agubosi, R. Oluwafemi, and J. O. Alagbe, 'The effect of processing on the proximate, mineral and vitamin composition of *Neem* leaves (*Azadirachta indica*) grown in Gwagwalada, FCT, Abuja', *Abuja Journal of Agriculture and Environment*, vol. 1, no. 1, pp. 293-299, 2021.
- [12] A. J. Olujimi, R. Sadiq Muhammad, M. Shittu, O. Daniel, and C. Olayemi, 'Effect of *Trichilia monadelpha* stem bark extract on the fatty acid composition of rabbit's thigh meat', *Journal of Environmental Issues and Climate Change*, vol. 1, no. 1, pp. 63-71, 2022. doi.org/10.59110/jeicc.v1i1.76
- [13] J. Alagbe, 'Use of medicinal plants as a panacea to poultry production and food security: A review', *Gospodarka I Innowacje*, vol. 22, pp. 1-12, 2022.
- [14] S. Sharma, S. Liu Xing, and K. Ram, 'Comparative analysis of ethanolic *Juniperus thurifera* leaf, stem bark and root extract using gas chromatography and mass spectroemetry', *International Journal of Agriculture and Animal Production*, vol. 2, no. 6, pp. 18-27, 2022. doi.org/10.55529/ijaap.26.18.27
- [15] R. A. Oluwafemi, O. C. Agubosi, and J. O. Alagbe, 'Proximate, minerals, vitamins and amino acid composition of *Prosopis africana* (African mesquite) seed oil', *Asian Journal of Advances in Research*, vol. 11, no. 1, pp. 21-27, 2021.
- [16] J. O. Alagbe, O. M. Adedeji, and D. Bamigboye, 'Influence of *Juniperus thurifera* root extract on the nutrient digestibility and caecal microbial count of growing rabbits', *Web of Synergy: International Interdisciplinary Research Journal*, vol. 1, no. 1, pp. 5-17, 2022.
- [17] J. O. Alagbe, '*Prosopis africana* (African mesquite) oil as an alternative to antibiotic feed additives on broiler chickens diets: haematology and serum biochemical indices', *Central Asian Journal of Theoretical and Applied Sciences*, vol. 3, no. 2, pp. 19-29, 2022.
- [18] J. O. Alagbe, '*Prosopis africana* (African mesquite) oil as an alternative to antibiotic feed additives on broiler chickens diets: performance and nutrient retention', *Discovery*, vol. 58, no. 314, pp. 134-142, 2022.
- [19] J. Alagbe, 'Chemical evaluation of proximate, vitamin and amino acid profile of leaf, stem bark and roots of *Indigofera tinctoria*', *International Journal on Integrated Education*, vol. 3, no. 10, pp. 150-157, 2020. doi.org/10.31149/ijie.v3i10.708
- [20] S. Peter, D. Stephene, and P. Eric, 'Unusal amount of (-) mesquitol from the heartwood of *Prosopis juliflora*', *Nat. Prod. Res*, vol. 23, pp. 183-189, 2009. doi.org/10.1080/14786410801940968
- [21] J. O. Alagbe, M. Shittu, and E. A. Ojo, 'Prospect of leaf extracts on the performance and blood profile of monogastric - A review', *International Journal of Integrated Education*, vol. 3, no. 7, pp. 122-127, 2020. doi.org/10.31149/ijie.v3i7.509

- [22] J. T. Barminas and H. Maina, 'Nutrient content of *Prosopis africana* seeds', *Journal of Plant Foods for Human Nutrition*, vol. 52, no. 4, pp. 325-328, 1998. <https://doi.org/10.1023/A:1008045218320>
- [23] E. Ankyambe-Akue, E. O. Ogunwolu, and R. Shaato, Potentials of Iron tree (*Prosopis africana*, Guill and Perr.) as protectant of cowpea seed (*Vigna unguiculata*) against the cowpea walp Bruchid (*Callosobruch maculatus* F). A paper read at Nigerian Society for Plant Protection 31st Annual Conference. Makurdi, 2004, pp. 22-26.
- [24] A. A. O. Ogunshe, M. O. Omotosho, and A. D. V. Ayanshina, 'Microbial studies and biochemical characteristics of controlled fermented afiyo- a Nigerian fermented food condiment from *Prosopis africana* (Guill and Perr)', *Taub.Pakistan Journal of Nutrition*, vol. 6, no. 6, pp. 620-627, 2007. doi.org/10.3923/pjn.2007.620.627
- [25] [Online]. Available: <http://www.fao.org/ag/agp/agpc/doc/gbase/data/pf00>. [Accessed: 20-May-2026].
- [26] J. O. Alagbe, O. C. P. Agubosi, R. Oluwafemi, and G. Zakara, 'Efficacy of *Trichilia monadelpha* stem bark on the growth performance of growing rabbits', *Best Journal of Innovation in Science*, vol. 1, no. 2, pp. 10-19, 2022. <https://doi.org/10.58489/2836-2217/005>
- [27] G. M. Gberikon, C. Agbulu, and M. E. Yaji, 'Nutritional composition of fermented powdered *Prosopis africana* soup condiment with or without inocula', *International Journal of Current Microbiology and Applied Sciences*, vol. 4, no. 2, pp. 166-171, 2015.
- [28] B. S. Platt, 'Biological Ennoblement; Improvement of the nutritive value of foods and dietary regimens by biological agencies', *Journal of Food Technology*, vol. 68, 1980.
- [29] F. A. Oguntoyinbo et al., 'Diversity of *Bacillus* species isolated from Okpepe, a traditional fermented soup condiment from Nigeria', *J. Food*, vol. 73, no. 5, 2010. doi.org/10.4315/0362-028X-73.5.870
- [30] A. O. Adewale, J. O. Alagbe, and A. O. Adeoye, 'Dietary Supplementation of *Rauvolfia Vomitoria* Root Extract as A Phytogetic Feed Additive in Growing Rabbit Diets: Haematology and serum biochemical indices', *International Journal of Orange Technologies*, vol. 3, no. 3, pp. 1-12, 2021.
- [31] J. Alagbe, 'Chemical evaluation of proximate, vitamin and amino acid profile of leaf, stem bark and roots of *Indigofera tinctoria*', *International Journal on Integrated Education*, vol. 3, no. 10, pp. 150-157, 2020. doi.org/10.31149/ijie.v3i10.708
- [32] O. Balogun, *The Federal Capital Territory of Nigeria. A Geography of its Development*. University of Ibadan Press Limited, 2001.
- [33] A. S. Singh, J. O. Alagbe, S. Sharma, R. Oluwafemi, and O. C. P. Agubosi, 'Effect of dietary supplementation of melon (*Citrullus linatus*) seed oil on the growth performance and antioxidant status of growing rabbits', *Journal of Multidimensional Research and Reviews*, vol. 2, no. 1, pp. 78-95, 2021. doi.org/10.47540/ijias.v1i2.175
- [34] D. Muritala, J. O. Alagbe, O. O. Ojebiyi, T. Ojadiran, and T. A. Rafiu, 'Growth performance and haematological and serum biochemical parameters of broiler chickens given varied concentrations of *Polyalthia longifolia* leaf extract in place of conventional antibiotics', *Animal Science and Genetics*, vol. 18, no. 2, pp. 57-71, 2022. doi.org/10.5604/01.3001.0015.9185
- [35] T. S. C. Silva, I. B. Suffredini, and E. L. Ricci, 'Antinociceptive and anti-inflammatory effects of *Lantana camara* L. extract in mice', *Rev Bras Plantas Med*, vol. 17, pp. 224-229, 2015. doi.org/10.1590/1983-084X/11109
- [36] H. Nakano, Y. Fujii, K. Yamada, S. Kosemura, S. Yamamura, and T. Suzuki, 'Isolation and identification of plant growth inhibitors as candidates for allelopathic substances from aqueous leachate from mesquite leaves', *Plant Growth Regulation*, vol. 37, pp. 113-117, 2002. doi.org/10.1023/A:1020579101938
- [37] J. O. Alagbe, S. Bamigboye, G. C. Nwosu, and D. Agbonika, 'Characterization of bioactive compounds in *Luffa aegyptiaca* leaf ethanolic extracts using gas chromatography and mass spectrometry (GC-MS)', *Drug Discovery*, vol. 17, pp. e10-1011, 2023. doi.org/10.54905/disssi.v17i39.e10dd1011
- [38] J. O. Alagbe, 'Bioactive compounds in ethanolic extract of *Strychnos innocua* root using gas chromatography and mass spectrometry (GC-MS)', *Drug Discovery*, vol. 17, pp. e4-1005, 2023. doi.org/10.54905/disssi.v17i39.e4dd1005

- [39] S. Velavan, 'Determination of phytochemicals in methanolic extract of *Annona muricata* leaf using GC-MS technique', *International Journal of Pharmacognosy Phytochemical Research*, vol. 7, no. 6, pp. 1251-1255, 2015.
- [40] B. Ti and F. O. Omotayo, 'Phytochemical and ethnobotanical study of some selected medicinal plants from Nigeria', *Journal of Medicinal Plants Research*, vol. 6, pp. 1106-1118, 2012. doi.org/10.5897/JMPR09.430
- [41] G. A. Ayoola, H. A. Coker, and S. A. Adesegun, 'Phytochemical screening and antioxidant activities of some selected medicinal plants used for malaria therapy in Southwestern Nigeria', *Tropical Journal of Pharmaceutical Research*, vol. 7, pp. 1019-1024, 2008. doi.org/10.4314/tjpr.v7i3.14686
- [42] K. S. Lokesh, D. Sonal, and M. P. Dobhai, 'Phytochemicals from *Prosopis* species', *Science and Technology*, vol. 4, no. 1, pp. 9-32, 2018.
- [43] S. Aneela and S. De, 'GC-MS analysis of methanolic extract of *Prosopis spicigera*', *International Journal of Phytopharmacology*, vol. 5, pp. 168-171, 2014.
- [44] S. Idris, G. I. Ndukwe, and C. E. Gimba, 'Preliminary phytochemical screening and antimicrobial activity of seed extracts of *Persea americana* (avocado pear)', *BAJOPAS*, vol. 2, pp. 173-176, 2009. doi.org/10.4314/bajopas.v2i1.58538
- [45] S. Valli, S. Gokulshankar, B. K. Mohanthy, M. S. Ranjith, S. Ashutosh, and V. Remya, 'Anticryptococcal activity of alkaloid rich fraction of leaves of *Prosopis juliflora*. A future promising supplementary therapy for cryptococcosis and cyyptococcal meningitis', *International Journal of Pharmaceutical Science*, vol. 6, pp. 491-495, 2014.
- [46] S. Sharma, S. Liu Xing, and K. Ram, 'Comparative analysis of ethanolic *Juniperus thurifera* leaf, stem bark and root extract using gas chromatography and mass spectroemetry', *International Journal of Agriculture and Animal Production*, vol. 2, no. 6, pp. 18-27, 2022. doi.org/10.55529/ijaap.26.18.27
- [47] V. K. Dua, A. C. Pandey, and A. P. Dash, 'Aducticidal activity of essential oil of *Lantana camara* leaves against mosquitoes', *Indian Journal Medicinal Research*, vol. 131, pp. 434-439.
- [48] C. U. Aguoru, C. Pilla, and J. O. Olasan, 'Phytochemical screening of *Xylopi aethiopia* with emphasis on its medicinally active principles', *Journal of Medicinal Plant Research*, vol. 7, pp. 1-5, 2016.
- [49] S. C. Okereke, U. Arunsi, and C. I. Nosiri, 'GC-MS/FT-IR screening of *Xylopi aethiopia* (Dunal) A. rich fruit', *African Journal Biochemistry Research*, vol. 11, no. 3, pp. 12-17, 2017.
- [50] L. J. Ogbadu, 'Studies on glutamic acid production by *Bacillus* isolates involved in daddawa fermentation', *Zaria. Pp*, vol. 1, 1988.

How to Cite Alagbe, John Olujimi. (2023). Bioactive compounds of prosopis africana oil (african mesquite) using gas chromatography and mass spectrometry (gc-ms) technique. *International Journal of Agriculture and Animal Production (IJAAP)*, 3(1), 89-97. <https://doi.org/10.55529/ijaap.33.1.10>

BIOGRAPHIE OF AUTHOR



Alagbe, John Olujimi, Ph.D., is a researcher and lecturer in the Department of Animal Nutrition and Biochemistry at Sumitra Research Institute, Gujarat, India. His research interests span feed biotechnology, animal toxicology, monogastric nutrition, plant extracts, and animal nutrition and management. He has made significant contributions to the field of phytogenic feed additives, focusing on the use of medicinal plant extracts and essential oils as natural alternatives to synthetic antibiotics in livestock and poultry production. His work has been cited over 2,400 times, reflecting his impact in the field. He collaborates extensively with researchers from Nigerian universities, including the University of Abuja and Ladoke Akintola University of Technology. Email: dralagbe@outlook.com