

Research Paper



Determining the effect of fe3o4 conjugated with chitosan nanoparticles on labneh product characteristics and its effect on blood picture parameters in anemic rats induced by phenylhydrazine

Hossam R. Attia^{1*}, Karakz M. Thalij²

^{1,2}Department of Food Sciences, College of Agriculture, Tikrit University, Tikrit, Iraq.

Article Info

Article History:

Received: 15 June 2023

Revised: 25 August 2023

Accepted: 01 September 2023

Published: 17 October 2023

Keywords:

Fe3O4

Anemia

Chitosan

Hb

Iron

Transferrin



ABSTRACT

The research aims to investigate the effect of Fe₃O₄-NPs and its conjugated with chitosan nanoparticles on the immunological indicators, such as IgA, IgG, and interleukin-6, and the iron indicators, of Ferritin, Transferrin, TIBC, and Iron in the serum of male albino rats induced with anemia by phenyl-hydrazine. The rats were separated into six groups in five replicates, each of which included (T1) the control group, (T2) anemia induced group, T3 group animals given Labneh enrichment with 2.5 mg of Fe₃O₄-NPs ml/day, T4 group animals given Labneh enrichment with 2.5 mg of Ch-NPs ml/day, T5 group animals given Labneh enrichment with 2.5 mg of each Fe₃O₄-NPs conjugated with Ch-NPs ml/day, T6 group animals given Labneh enrichment with 2.5 mg of Hamvir ml/day, and were fed for 30 days. The results showed a change and increase in the values of free fatty acids, as they reached their highest value after 20 days of preservation, where the treatment (T1) recorded a noticeable increase, reaching (2.31). Likewise, in the treatment (T5), we notice a significant increase in it, as it reached (0.75), in contrast to the rest. All coefficients that were (T2), (T3) and (T4) recorded (0.53), (0.58) and (0.63), respectively. the values of RBCs and Hb, which gave a significant decrease and it was at 2.84 (1×10⁶/ ml) and 6.4 (g/dl), respectively, compared to the control treatment (T1).

Corresponding Author:

Hossam R. Attia

Department of Food Sciences, College of Agriculture, Tikrit University, Tikrit, Iraq.

Email: hram.alward@gmail.com

Copyright © 2023 The Author(s). This is an open access article distributed under the Creative Commons Attribution License, (<http://creativecommons.org/licenses/by/4.0/>) which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

1. INTRODUCTION

The dairy product labneh, which is made by fermenting milk with lactic acid bacteria to give it a distinctive flavour and scent, is very well-liked throughout the world, but especially in Middle Eastern countries reducing lactose intolerance symptoms, serving as a significant calcium source, and boosting the number of good bacteria in the intestines. Labneh is a useful food and a good source of essential nutrients. [1].

On the other hand, many countries adopted nanotechnology in the treatment of many diseases, including incurable ones, as it was noted for its ability to cure some diseases and is used daily in hospitals and health centers to preserve human health through early detection of diseases, which reduces the cost of treatment [2]. Fe₃O₄-NPs are widely used in numerous significant biological and medicinal applications, and because they are super magnetic, they have various uses in sophisticated technology. [3] It is used in drug delivery and loading systems, as it has the property of carrying more than one treatment material at one time, which includes many diseases, including bacterial and fungal diseases [4]. It also has several applications, including adsorption applications and heavy water treatment applications [5]. Chitosan results from the removal of acetyl groups from chitin, which is considered the second most common compound in nature after cellulose, and its molecular formula is (C₆H₁₁O₄N)_n, which constitutes (20-30)% of crustacean shells such as shrimp and crabs [6] As for nanoscale chitosan (Ch-NP) It has the properties of chitosan and properties of nanoparticles such as the effect of surface area and small size [7]. Chitosan nanoparticles (ChNP) were first described in 1994 by (Ohya) and colleagues. They used ChNP prepared by emulsification and crosslinking to deliver the anti-cancer drug 5-fluorouracil intravenously [8] On the other hand, anemia is viewed as a global public health issue caused by a shortage of red blood cells or a reduction in their ability to carry enough oxygen to meet the needs of the physiological body. The severity of anemia varies with age group, gender and during pregnancy, in 2015, the World Health Organization (WHO) estimates that about 1.62 billion (24.8%) of the world's population suffer from anemia, with a higher rate among pregnant women, as it is The proportion of anemia cases is 38% (32.4 million), 29% (496 million) are non-pregnant, and 43% (273 million) are in children. Anemia may have multiple causes, but according to WHO (2015), 50% of cases of anemia blood caused by an insufficient amount of iron [9]. Due to the lack of studies on the effect of laboratory-made labneh fortified with Fe₃O₄-NPs, and the complex of chitosan with Fe₃O₄-NPs Effects on Complete Blood Picture.

2. METHODOLOGY

Preparation of the Fe₃O₄-NPs with Ch-NPs: 10 mg of Fe₃O₄-NPs was added to 1g of Ch-NPs (NANOSHEL, USA) in 10 ml of deionized aqueous solution and stirred at room temperature for 24 hours. The required amount of the conjugated nanoparticles compound (CNC) was dried using a lyophilized technique and kept at 4 °C in the refrigerator until use.

Labneh Preparation: the preparation of the support tray template with nanomaterials and the drug Hemafer: local cow's milk was obtained from Salah Al-Alin province (Al-Alam district), and the brick was manufactured according to the method mentioned before [10]. Where the milk is heated to 85 °C for half an hour, then the milk is cooled to 42 °C, and the traditional starter culture is added at a rate of 3%. Distributing the salt, then the filtering process is carried out with bags made of cloth type (Japan), which are placed suspended in the refrigerator for 20 hours, after which they were fortified with the aforementioned nanomaterials and the drug at a rate of 10 mg/100 gm of labneh.

Determination of free Fatty Acids: Free fatty acids were examined according to what was mentioned in [11], by mixing 30 ml of diethyl ether solvent with 30 ml of ethanol alcohol with 1 ml of phenolphthalein indicator. The previous mixture was neutralized using 0.1 sodium hydroxide until the pink color appeared. Then put 10 grams of labneh in the neutralizing solution, mix well, then filter the solution, and the filtrate is calibrated with the previous rule until the neutralization point is reached, which is identified by the appearance of a pink color in the solution. The amount of base consumed after adding the labneh was

calculated, through which the amount of free fatty acids was calculated, which is expressed as a percentage of free fatty acids in the labneh and is calculated as oleic acid. According to the following equation:

$$\text{FFA} = 2.82 \times \text{T/W}$$

PH Measurement: 10 g of labneh was weighed and digested well in a ceramic mortar with 10 ml of distilled water and a pH-meter was used to estimate the pH value of the sample [12].

2.1. Initialization of Laboratory Animals

Albino Sprague-Dawley weanling, Male laboratory rats, were obtained from the College of Veterinary Medicine-Tikrit University, at the age of 8-9 weeks. Their weights ranged between 150-152 g, and they were separated randomly into groups of similar weights.

The animals were placed in cages made of plastic, and the floor was covered with sawdust, which was replaced four times a week. The animals were fed regularly using the ready-made diet according to what was mentioned in [13], and the basic food contained (g / kg diet) two cases: 84.95 g, net protein: 158.5 g, oil: 100 g, and a mixture of vitamins: 5 g and 50 g. A mixture of mineral salts, 50 gm of cellulose, 100 gm of glucose, and 536.5 gm of starch. The animals were raised under the supervision of specialized veterinary staff, taking into account the aspect of hygiene.

Experimental animals were distributed into six groups each one including five animals, that included: T1 healthy control group (without treatment, which was given water and food only for the duration of the experiment), T2 induced anemia infection group (given 40 mg by subcutaneous intraperitoneal of phenyl hydrazine/kg of body weight each day for 48 hours) [14], T3 group animals given Labneh enrichment with 2.5 mg of Fe₃O₄-NPs ml/day, T4 group animals given Labneh enrichment with 2.5 mg of Ch-NPs ml/day, T5 group animals given Labneh enrichment with 2.5 mg of each Fe₃O₄-NPs conjugated with Ch-NPs ml/day, T6 group animals given Labneh enrichment with 2.5 mg of Hamvir ml/day. Diets and water were free of ad lantum given, and 0.1 mg of phenyl hydrazine was given for each kg of body weight animal by subcutaneous intraperitoneal throughout the experiment.

2.2. Stages of the Experiment

The phases of the vital experiment included the following:

The First Stage (pre-treatment stage with anemia): Animals in groups (T1-TG6) were fed the usual diet for 5 days, after which blood samples were drawn from each group in two tubes, the first containing EDTA anticoagulant to measure complete blood picture, and the other free of it containing Approximately 2-3 ml of blood was centrifuged using a centrifuge at a speed of 3000 cycles/ min for 15 minutes to obtain serum that was kept at -20 ° C until analyzes were performed, as in [15], to adopt them as control samples before induced anemia.

The Second Stage (treatment with immunosuppression): animals in groups (T2, T3, T4, T5 and T6) were treated with 40 mg by subcutaneous intraperitoneal of phenyl hydrazine/kg of body weight each day for 48 hours until the appearance of symptoms of anemia, which included lethargy, lack of movement, withdrawal and poor appetite. for food that appeared after 3 days. Blood samples were drawn from the groups in two tubes, as in the above paragraph, to ensure that anemia occurred compared to the samples of the first stage.

The Third Stage (treatment with each of Fe₃O₄-NPs or Ch-NPs singly or in conjugated): induced anemia animals in groups (T3, T4, T5 and T6) were treated with each of the Fe₃O₄-NPs or Ch-NPs singly or in conjugated. After the end of the experiment, blood samples were drawn as in the two paragraphs above.

Measured of Biochemicals Parameters: The CBC Mindray device was used according to the manufacturer's instructions, and the blood tests included estimating the hemoglobin concentration (Hb), the red blood cell count (RBCs), the total white blood cell count (WBC), measuring the percentage of packed blood cell volume (P.C.V.) packed cell volume and the average red cell volume. (MCV) and platelet count.

Statistical Analysis: The biological experiment was implemented according to the randomized complete design CRD, and the analysis of variance was carried out using the General Linear Model within the [16] SAS program. When there are significant differences between the means, Duncan's test [17] was used to determine the significance of the differences between the different means at a probability level of 0.05.

3. RESULTS AND DISCUSSION

3.1. Determination of Free Fatty Acids (FFA) in Milk Manufactured and Treated with Nanomaterials

Estimating the values of free fatty acids in the labneh manufactured in the laboratory is evidence of its suitability during preservation and experiment. In Table 1 we note from the results that there are no significant differences ($P < 0.05$) at the initial time for all treatments whose rates were between (0.4) - (0.8) However, there are significant differences after 10 days of preservation, especially treatment (T1) the control when compared with other treatments, and the least change was (T2) and (T4) compared with other treatments, and the results showed a change and increase in the values of free fatty acids, as It reached its highest value after 20 days of preservation, as the transaction (T1) recorded a significant increase, amounting to (2.31), as well as the transaction (T5), we notice a significant increase in it, as it reached (0.75). In contrast to all the treatments that were (T2), (T3) and (T4), which recorded (0.53), (0.58) and (0.63), respectively, the reason for the decrease in the percentage of free fatty acids in these treatments can be attributed to the role of nanoparticles in inhibiting microorganisms and Gram-negative bacteria that contaminate brick, this is the first study in this field.

Table 1. Values of Free Fatty Acids (FFA) In Yoghurt Manufactured and Treated with Nanomaterials after Storage for 20 Days at a Temperature of $5^{\circ}\text{C} \pm 2$

Treatment Type	The Values of Free Fatty Acids in Labneh Samples Manufactured After A 20-Day Storage Period At $5^{\circ}\text{C} \pm 2$		
	First Hour	10Days	20 Days
T1	0.8a \pm 0.01	0.98a \pm 0.01	2.31a \pm 0.01
T2	0.4b \pm 0.02	0.43c \pm 0.02	0.53d \pm 0.02
T3	0.4b \pm 0.02	0.45bc \pm 0.01	0.58c \pm 0.01
T4	0.5b \pm 0.01	0.43c \pm 0.01	0.63c \pm 0.01
T5	0.5b \pm 0.01	0.51b \pm 0.02	0.75b \pm 0.02

The different letters in the same column indicate that there are significant differences T1: control, T2: nano-Fe₃O₄, T3: nano-chitosan, T4: chitosan loaded with nano-Fe₃O₄, T5: HEMAFER (Ferric hydroxide polymaltose complex).

3.2. Estimation of the PH in the Manufactured Brick Treated with Nanomaterials

The pH value indicates the natural and developed acidity, which usually indicates the presence of lactic acid bacteria and the effectiveness of its enzymes in fermenting lactose and converting it into lactic acid, where it participates with other compounds and gives taste and flavor, which gives an indication of its acceptability to the consumer. The results are shown in Table 2 the pH value was similar for all treatments at the beginning of manufacturing.

Table 2. PH Values in the Manufactured Brick Treated with Nanomaterials after Storage for 20 Days at $5^{\circ}\text{C} \pm 2$.

Treatment Type	The Values of PH in Labneh Samples Manufactured after A 20-Day Storage Period At $5^{\circ}\text{C} \pm 2$		
	first hour	10days	20 days
T1	4.72a \pm 0.4	4.10b \pm 0.6	3.05b \pm 0.5
T2	4.75a \pm 0.5	4.23a \pm 0.4	4.15a \pm 0.3
T3	4.74a \pm 0.3	4.46a \pm 0.7	4.24a \pm 0.6
T4	4.72a \pm 0.7	4.33a \pm 0.3	4.29a \pm 0.7
T5	4.69a \pm 0.2	4.15b \pm 0.4	4.01b \pm 0.5

The different letters in the same column indicate that there are significant differences T1: control, T2: nano-Fe₃O₄, T3: nano-chitosan, T4: chitosan loaded with nano-Fe₃O₄, T5: HEMAFER (Ferric hydroxide polymaltose complex).

That is, their rates were between 4.69 – 4.75. The pH in the first 10 days of storage for treatment (T1) and (T5) was at 4.10 and 4.15, respectively. As for the end of the experiment, that is, after 20 days of preservation, the decrease in the pH value continued for the treatments (T1) and (T5), as it was at 3.05 and 4.01, while the treatments to which the studied nanomaterials were added (T2), (T3) and (T4) were at 4.15, 4.24, and 4.29. The evolution in the pH value of the treatments (T1) and (T5) is attributed to the activity of the microorganisms present in the labneh product and thus the fermentation of the remaining lactose [18].

3.3. Effect of Oral Dosage with Fe₃O₄-NPs Conjugated with Ch-NPs on Physiological Parameters of Anemia-Induced Rats with phenyl-hydrazine.

Effects on Complete Blood Picture: The effect of oral dosage of different treatments on serum iron indicators of anemia-induced rats with phenylhydrazine was noted in Table 3 the impact of doses of nanoparticles, nano complex and hemavir on serum iron levels and indicators of anemia-induced animals.

Table 3. Effect of Oral Dosage with Fe₃O₄-Nps Conjugated with Ch-NPs on Blood Picture of Anemia-Induced Rats with Phenyl-Hydrazine

Treatment Type	Symbol	RBCs	Hb	HCT	M.C. V	Platelet
		(1×10 ⁶ /ml)	(g/dl)	%	μM ³	(10 ⁹ /L)
Control	T1	6.63a±0.8	12.8a±1.3	39.7b±3.8	59.6d±3.7	295.4d±21.5
Infection with anemia induced	T2	2.84d±0.2	6.4c±0.9	19.1d±2.4	67.3a±5.2	561.6a±20.8
Treatment	T3	4.85c±0.3	12.7a±1.8	40.4b±2.7	60.6d±4.6	319.3c±19.4
	T4	4.86c±0.3	12.3a±1.0	42.9a±3.5	63.6c±5.3	322.6c±15.7
	T5	5.27b±0.4	12.9a±1.2	41.1b±3.3	63.3c±5.1	314.6c±16.6
	T6	5.96b±0.5	11.3b±1.1	35.8c±2.9	65.6b±4.7	530.2b±21.5

3.4. The Different Letters in the Same Column Indicate that there are Significant Differences. T1: Control Group, T2: Group Orally phenyl-hydrazine Induced Anemia, T3: Group Orally with Fe₃O₄-NPs, T4: Orally with Ch-NPs, T5: Orally with Fe₃O₄-NPs Conjugated with Ch-NPs, T6: Orally with HEMAFER Drug

The effect of nanoparticles of Fe₃O₄ and nanoparticles of chitosan, in addition to the complex of chitosan with Fe₃O₄ nanoparticles, of rats dosed orally for 30 days, can be seen in the parameters of the blood picture, which were shown in Table (3). normal levels of their levels in the blood of laboratory rats dosed with the aforementioned substances, where the values of RBCs for treatments T3, T4, T5, and T6 gave 4.85, 4.86, 5.27, and 5.96 (1 x 10⁶/ml), respectively, in contrast to treatment T2, which gave a significant decrease when compared T1-treated control.

The results also showed that the value of the hemoglobin Hb concentration decreased significantly in the blood of the group of animals in which anemia was induced experimentally with PHZ and untreated (T2) and was at 6.56 (g / dl) when compared to the groups of treatments T3, T4, T5 and T6 that were treated with nanomaterials and the drug HEMAFER Significant good results, which were 12.7, 12.3, 12.9, and 11.3 (g/dl), respectively, compared with the control group (T1), which was valued at 12.8 (g/dl). We also note that the percentage of agglutinated blood cells (HCT) decreased in the group of animals (T2), when it was at 19.1%, compared with the group of healthy control animals (T1), which was 39.7%. While the results increased significantly after treatment in groups (T3), (T4), ((T5, and (T6), they were at 40.4, 42.9, 41.1, and 35.8%, respectively, compared with the treatment value (T2).

We also notice that the red blood cells (M.C.V) suffered a significant increase in the untreated (T2) group of animals, as it gave a value of 67.3 (μM³), while we notice a significant decrease in its values after treatment in the groups (T3), ((T4), and ((T5). At 60.6, 63.6, and 63.3 (μM³), respectively, compared with

(T2), but we find that (T6) remained slightly higher than its predecessors, giving 65.6 (μM^3) when compared with the value of the T1 control group, which gave a value of 59.6 (μM^3). As well as for platelets PLT, it recorded a significant decrease in the groups of treated animals (T3), (T4), and (T5) and it was at 319, 322 and 314 (103/L) respectively, compared with the two groups (T2) and (T6) where it was at 561 and 530 (103/L) respectively, while in the control group (T1) it was at 295 (103/L).

The results of the study in groups of animals treated with phenylhydrazine and its effects on various blood variables, which include the number of red blood cells, the size of compact cells, and Hb, which is the most important variable indicating anemia and some other variables, agreed with what was mentioned by [19]. The research indicated a decrease in the results of the values of these variables that were studied due to the toxicity of the drug phenylhydrazine resulting from oxidative stress that leads to the generation of free radicals, which in turn attack vital molecules, causing damage to the biological system [20].

These results agreed with the findings of [21], who indicated an improvement in the levels of RBCs, Hb and PCV concentrations and a decrease in M.C.V in laboratory rats when they were dosed with 1.5 mg of Fe₃O₄ NPs. [22] also mentioned that nano-chitosan has the ability to enhance the absorption of macro- and micronutrients, including iron. The results of the researcher, [23]. showed that fortifying the food with nano-chitosan led to an improvement in the levels of Hb, Hct, MCV, and the number of red blood cells in fish fed on it.

Acknowledgments

The authors have no specific acknowledgments to make for this research.

Funding Information

This research received no specific grant from any funding agency in the public, commercial, or not-for-profit sectors.

Author Contributions Statement

Name of Author	C	M	So	Va	Fo	I	R	D	O	E	Vi	Su	P	Fu
Hossam R. Attia	✓	✓	✓	✓		✓	✓	✓	✓	✓	✓		✓	✓
Karakz M. Thalij	✓	✓	✓		✓		✓	✓	✓		✓		✓	✓

C : Conceptualization

M : Methodology

So : Software

Va : Validation

Fo : Formal analysis

I : Investigation

R : Resources

D : Data Curation

O : Writing - Original Draft

E : Writing - Review & Editing

Vi : Visualization

Su : Supervision

P : Project administration

Fu : Funding acquisition

Conflict of Interest Statement

The authors declare that there are no conflicts of interest regarding the publication of this paper.

Informed Consent

All participants were informed about the purpose of the study, and their voluntary consent was obtained prior to data collection.

Ethical Approval

The study was conducted in compliance with the ethical principles outlined in the Declaration of Helsinki and approved by the relevant institutional authorities.

Data Availability

The data that support the findings of this study are available from the corresponding author upon reasonable request.




REFERENCES

- [1] E. J. Ibrahim, K. M. Thalij, M. K. Saleh, and A. S. Badawy, 'Biosynthesis of Zinc Oxide Nanoparticles and Assay of Antibacterial Activity', *American Journal of Biochemistry and Biotechnology*, vol. 13, no. 2, pp. 63-69, 2017. doi.org/10.3844/ajbbsp.2017.63.69
- [2] A. Fouda and T. I. Shaheen, 'Silver nanoparticles: biosynthesis, characterization and application on cotton fabrics', *Microbiology Research Journal International*, vol. 20, 2017. doi.org/10.9734/MRII/2017/32961
- [3] S. Laurent, S. Boutry, I. Mahieu, L. Vander Elst, and R. N. Muller, 'Iron oxide-based MR contrast agents: from chemistry to cell labelling', *Curr Med Chem*, vol. 16, 2009. doi.org/10.2174/092986709789878256
- [4] Arias, L. S. Pessan J. P. Ana Paula. M. V, Taynara M.T. de L. Alberto C. B. D and Douglas R. M, (2018). "Iron Oxide Nanoparticles for Biomedical Applications: A Perspective on Synthesis, Drugs, Antimicrobial Activity, and Toxicity" *Antibiotics* 10.3390/antibiotics.7020046 doi.org/10.3390/antibiotics7020046
- [5] A. M. Gutierrez, T. D. Dziubla, and J. Z. Hilt, 'Synthesis of superparamagnetic Fe₃O₄ Nanoparticles coated with green tea polyphenols and their use for removal of dye pollutant from aqueous solution"', *Journal of Environmental Chemical Engineering*, vol. 5, no. 3, 2017. doi.org/10.1016/j.jece.2017.04.022
- [6] M. Dash, F. Chiellini, R. M. Ottenbrite, and E. Chiellini, 'Chitosan-a versatile semi-synthetic polymer in biomedical applications', *Progress in Polymer Science*, vol. 36, no. 8, pp. 981-1014, 2011. doi.org/10.1016/j.progpolymsci.2011.02.001
- [7] A. Ingle, A. Gade, S. Pierrat, C. Sonnichsen, and M. Rai, 'Mycosynthesis of silver nanoparticles using the fungus *Fusarium acuminatum* and its activity against some human pathogenic bacteria', *Curr Nanosci*, vol. 4, no. 2, pp. 141-144, 2008. doi.org/10.2174/157341308784340804
- [8] A. Grenha, 'Chitosan nanoparticles: a survey of preparation methods', *J Drug Target*, vol. 20, no. 4, pp. 291-300, 2010. doi.org/10.3109/1061186X.2011.654121
- [9] G. A. Stevens, M. M. Finucan, L. M. De-Regil, C. J. Paciorek, S. R. Flaxman, and F. Branca, 'Global, regional, and national trends in haemoglobin concentration and prevalence of total and severe anaemia in children and pregnant and non-pregnant women for 1995-2011: a systematic analysis of population-representative data', *Lancet Glob Health*, vol. 1, no. 1, pp. e16-e25, 2013. [doi.org/10.1016/S2214-109X\(13\)70001-9](https://doi.org/10.1016/S2214-109X(13)70001-9)
- [10] Dietary Allowance. Washington. D.C: National Academy. Press, 2002.
- [11] 'Official Methods of Analysis of AOAC International', AOAC International. Ch. 17. USA, 2010.
- [12] R. Hool et al., Chemical and Physical Methods. In: *Standard Methods for the Examination of Dairy Products* Wehr. Washington: American Public Health Association, 2004, pp. 363-532.
- [13] A. Y. Tamime, Science and Technology. Oxford, UK: Pergamon Press Ltd, 1985.
- [14] H. Anbara, R. Shahrooz, M. Razi, H. Malekinejad, and G. Najafi, 'The effect of vitamin C on mice hemolytic anaemia induced by phenylhydrazine: an animal model study using histological changes in testis, pre-implantation embryo development, and biochemical changes', *Iranian journal of basic medical sciences*, vol. 21, no. 7, 2018.
- [15] Y. Tietz, *Clinical Biochemistry*; 6th ed, vol. 825. New York: McGraw-Hill, 2005.
- [16] Duncan, D.B. (1955). Multiple ranges and multiple "F" tests. *Biometric*, 11:1-42. doi.org/10.2307/3001478
- [17] 'Statistical Analysis System, User's Guide. Statistical. Version 9', SAS. Inst. Inc. Cary. N.C. The USA, 2012.
- [18] I. Fathollahi, J. Hesari, S. Azadmard, and S. Oustan, 'Influence of Proteolysis and Soluble Calcium Levels on Textural Changes in the Interior and Exterior of Iranian UF White Cheese during Ripening', *World Acad. Sci. Eng. Technol*, vol. 66, pp. 844-849, 2010.
- [19] M. E. Shahenda and B. A. Jehan, 'The anti-anemic effect of dried beet green in phenylhydrazine treated rats', *Arch Pharm Sci ASU*, vol. 2, no. 2, pp. 54-69, 2018. doi.org/10.21608/aps.2018.18735

- [20] S. Sharma and C. Haldar, 'Comparative effect of melatonin and vitamin E on phenylhydrazine-induced toxicity in the spleen of *Funambulus pennanti*', *Environmental Toxicology: An International Journal*, vol. 24, no. 1, pp. 1-9, 2009. doi.org/10.1002/tox.20383
- [21] R. Sobeh, H. M. Amin, S. E. Saleh, and A. Souad, 'Evaluation of the radio-protective role of PEG Fe₃O₄ NPs on γ -irradiated male Wistar rats', *Environmental Nanotechnology, Monitoring & Management*, vol. 17, 2022. doi.org/10.1016/j.enmm.2021.100620
- [22] F. S. Abd El-Naby, M. A. E. Naiel, A. A. Al-Sagheer, and S. S. Negm, 'Dietary chitosan nanoparticles enhance the growth, production performance, and immunity in *Oreochromis niloticus*', *Aquaculture*, vol. 501, pp. 82-89, Feb. 2019. doi.org/10.1016/j.aquaculture.2018.11.014
- [23] M. El-Naggar, H. El-Shabaka, M. Khalil, and A. Suloma., Eds, Efficacy of dietary chitosan and chitosan nanoparticles supplementation on health status of Nile tilapia, *Oreochromis niloticus*. Cairo; Egypt, 2021. doi.org/10.1016/j.aqrep.2021.100628

How to Cite Hossam R. Attia, Karakz M. Thalij. (2023). Determining the effect of fe₃o₄ conjugated with chitosan nanoparticles on labneh product characteristics and its effect on blood picture parameters in anemic rats induced by phenylhydrazine. *International Journal of Agriculture and Animal Production (IJAAP)*, 3(2), 92-99. <https://doi.org/10.55529/ijaap.36.1.9>

BIOGRAPHIES OF AUTHORS

	<p>Hossam R. Attia, is a prominent academic and food scientist affiliated with the Department of Food Sciences, College of Agriculture at Tikrit University, Tikrit, Iraq. With specialized expertise in food sciences and agricultural research, he has made significant contributions to advancing knowledge in food technology, safety, and quality within Iraq's academic sphere. His research endeavors at Tikrit University's College of Agriculture reflect a deep commitment to addressing food security challenges and promoting sustainable agricultural practices. Hossam's scholarly work positions him as a valuable contributor to Iraq's growing food science research community and broader agricultural development initiatives. Email: hram.alward@gmail.com</p>
	<p>Karakz M. Thalij , is an emerging and dedicated researcher in the field of food sciences, associated with the Department of Food Sciences, College of Agriculture at Tikrit University, Tikrit, Iraq. Her academic pursuits are centered on exploring innovative approaches to food technology, nutrition, and agricultural science, contributing meaningfully to Iraq's food research landscape. With a strong passion for scientific inquiry and problem-solving, she actively engages in research that addresses critical food quality and safety challenges facing Iraqi communities. Karakz's commitment to excellence at Tikrit University reflects her determination to make a lasting impact on food science advancement in the region. Email: kthalij@tu.edu.iq</p>