International Journal of Research in Science & Engineering
ISSN: 2394-8299
Vol: 02, No. 02, Feb-Mar 2022
http://journal.hmjournals.com/index.php/IJRISE
DOI: https://doi.org/10.55529/ijrise.22.29.40

Physics of Cooking: Heat Transfer and Nutritional Retention

O. C. Molua[1], A. O. Ukpene[2*], Emagbetere. J. U[3], C. P. Ukpene[4], Apaokwueze. T.N[5]

[1]Physics Department, University of Delta, Agbor, Nigeria.
[2]Biological Sciences Department, University of Delta, Agbor, Nigeria.*
[3]Physics Department, College of Education, Mosogar, Nigeria.
[4,5]Home Economics Department, University of Delta, Agbor, Nigeria.

Email: [1]collins.molua@unidel.edu.ng, [3]johnbullemagbetere@yahoo.com,
[4]patience.ukpene@unidel.edu.ng, [5]tessy.apaokwueze@unidel.edu.ng
Corresponding Email: [2]anthony.ukpene@unidel.edu.ng*

Received: 16 November 2021 **Accepted:** 04 February 2022 **Published:** 20 March 2022

Abstract: ***This article examines the field of culinary physics, investigating the complex correlation between heat transfer in cooking procedures and its influence on the nutritional content of food. By examining the mechanisms underlying heat transfer and their impact on the preservation of nutritional content, this study can bring about a transformative shift in culinary techniques. By attaining a state of balance between energy efficiency, sensory appeal, and the retention of essential nutrients, this research can revolutionize the field of cooking. This research endeavor's amalgamation of physics, home economics, and biology presents opportunities for a more profound comprehension of the culinary arts.***

Keywords: Culinary Physics, Heat Transfer, Nutritional Retention, Cooking Methods.

1. INTRODUCTION

Cooking is a time-honored tradition that converts unprocessed components into delectable and nutritious culinary creations. The culinary process encompasses more than a mere amalgamation of ingredients; it involves a nuanced interaction of physical and chemical transformations governed by the principles of thermal energy transfer [1] [2]. The transfer of heat during the cooking process has a notable impact on the nutritional composition of food, leading to a multifaceted area of study that combines principles from physics, home economics, and biology [3]. Cooking, a form of artistic expression that has undergone significant transformations over centuries, is an essential and universal human activity that surpasses cultural barriers. The process converts unprocessed components into appetizing

Copyright The Author(s) 2022.This is an Open Access Article distributed under the CC BY license. (http://creativecommons.org/licenses/by/4.0/)

International Journal of Research in Science & Engineering
ISSN: 2394-8299
Vol: 02, No. 02, Feb-Mar 2022
http://journal.hmjournals.com/index.php/IJRISE
DOI: https://doi.org/10.55529/ijrise.22.29.40

culinary creations, each possessing a unique taste profile and nutritional makeup [4]. In addition to its cultural significance, cooking is firmly grounded in physics principles, specifically the complex heat transfer mechanisms. How heat is distributed and absorbed during the culinary process significantly impacts the texture, taste, and nutritional composition of food [5].

The field of culinary physics encompasses a diverse range of scientific principles integrated with the artistic aspects of cooking. Each culinary technique, ranging from sautéing to roasting, can be understood as an embodiment of heat transfer mechanisms in operation. Heat transfer from the heat source to the food initiates a series of physical and chemical transformations that give rise to the appealing textures and flavors commonly associated with prepared dishes [6]. In addition to sensory attributes such as taste and texture, a comprehensive comprehension of the principles of heat transfer in the context of cooking reveals its profound influence on the nutritional composition of food [7]. This aspect of culinary science holds significant relevance for disciplines such as home economics and biology.

Literature Reviews

Prior research has demonstrated the significant impact of heat transfer on the preservation of nutrients during cooking. The rate and extent of nutrient loss are significantly influenced by heat transfer mechanisms, namely conduction, convection, and radiation [8]. For example, the blanching technique, which encompasses the application of swift heating and subsequent cooling, has demonstrated superior efficacy in maintaining vegetables' color, flavor, and nutritional content when contrasted with conventional boiling methods [9] . Upon delving into the existing body of literature, one discovers a vast array of scholarly research that highlights the fundamental significance of heat transfer in the culinary realm and its profound implications for preserving nutritional value. The process of conduction, which involves the transfer of heat through direct contact between a heat source and the food, has been the subject of extensive research [10] [11] [12]. The phenomenon mentioned above is prominently observable in culinary methods such as grilling and pan-frying, wherein the direct contact between the food and the heated surface induces the Maillard reaction. This intricate chemical process intensifies taste and modifies the nutritional makeup.

Convection, as an additional mechanism, encompasses the process of heat transfer through the motion of fluids, which can be in the form of either liquids or gases. In the culinary domain, convection is a phenomenon that manifests in various cooking techniques, such as boiling and baking [13]. The process of boiling vegetables can result in the loss of nutrients, such as vitamins, as they are released into the water used for cooking. Nevertheless, recent research has indicated that reducing the water-to-vegetable ratio and incorporating the cooking liquid into the consumption process can aid in preserving certain nutrients that would otherwise be lost.

Radiation, an often overlooked mechanism, encompasses the transfer of heat through electromagnetic waves, specifically infrared radiation, which is commonly employed in grilling and baking processes [14][15]. This approach has the potential to facilitate expeditious cooking and the formation of a browned crust on the surface of the food. The process of browning in food preparation contributes to enhanced flavor and texture, but it can

Copyright The Author(s) 2022.This is an Open Access Article distributed under the CC BY license. (http://creativecommons.org/licenses/by/4.0/)

International Journal of Research in Science & Engineering
ISSN: 2394-8299
Vol: 02, No. 02, Feb-Mar 2022
http://journal.hmjournals.com/index.php/IJRISE
DOI: https://doi.org/10.55529/ijrise.22.29.40

also give rise to the creation of potentially deleterious compounds [16]. Evaluating the advantages and drawbacks associated with radiation-based cooking techniques continues to be a subject of scholarly inquiry.

In order to examine the complex relationship between heat transfer and nutritional retention, a comprehensive experimental methodology is employed. This study investigates using diverse cooking techniques, including boiling, steaming, frying, and baking, in conjunction with various ingredients. Temperature probes are utilized to monitor the distribution of heat within food samples, while nutritional analyses are performed before and after the cooking process. This enables a quantitative evaluation of nutrient degradation and its association with heat transfer mechanisms. In order to elucidate the complex interplay between heat transfer and nutritional retention, a comprehensive and systematic methodology is utilized. A variety of cooking techniques and ingredients are selected as variables for investigation. The techniques employed include boiling, steaming, frying, and baking, while the ingredients include vegetables, meats, and grains. The observation of each method is conducted with great attention to detail, employing a combination of temperature probes, thermal imaging, and nutrient analysis techniques.

2. METHODOLOGY

Before the commencement of the cooking process, a comprehensive nutritional analysis is conducted on the raw ingredients to establish a foundational reference point. Following the cooking process, the samples undergo identical analysis to quantify the degree of nutrient degradation. The methodical nature of this approach facilitates a comprehensive comprehension of the specific effects of each heat transfer mechanism on various nutrients.
The methodology encompasses investigating novel culinary methods that leverage principles of physics to maximize the preservation of nutrients. One example of a cooking technique that has gained popularity is sous-vide cooking. This method utilizes precise temperature control within a water bath to effectively preserve flavors and nutrients. This methodological framework sheds light on the complex relationship between physics, culinary arts, and nutritional science, providing valuable insights that have the potential to revolutionize our approach to cooking in terms of both flavor and health considerations.

In order to gain a comprehensive understanding of the physics involved in the cooking process, as well as the various mechanisms of heat transfer and their impact on the retention of nutritional components, a methodical approach was employed.

The initial phase of the study entailed establishing the research parameters and developing specific research goals. This entailed investigating various cooking techniques, evaluating the efficacy of heat transfer, assessing the preservation of nutrients, examining the impact of cooking duration, considering different food categories, and analyzing the influence of temperature on nutrient retention.

The second phase entailed a comprehensive examination of scholarly literature, research papers, studies, and articles on the physics of culinary processes, heat transfer phenomena, and alterations in nutritional composition. This study facilitated comprehension of extant research about diverse cooking techniques and their influence on nutritional composition.

Copyright The Author(s) 2022.This is an Open Access Article distributed under the CC BY license. (http://creativecommons.org/licenses/by/4.0/)

International Journal of Research in Science & Engineering
ISSN: 2394-8299
Vol: 02, No. 02, Feb-Mar 2022
http://journal.hmjournals.com/index.php/IJRISE
DOI: https://doi.org/10.55529/ijrise.22.29.40

The third step involved the analysis of various cooking methods, focusing on evaluating their efficiency and mechanisms of heat transfer. This involved comprehending the mechanisms by which heat is transferred in culinary procedures, including conduction, convection, and radiation.

The fourth step entailed investigating the impact of various cooking techniques on the preservation of nutrients and analyzing the effects of these methods on vitamins, minerals, and other essential nutritional elements.

The fifth step of the process involved examining the relationship between cooking time and the preservation of nutrients. This study examined the impact of prolonged cooking times on the nutritional composition of food items.

The sixth step entailed investigating the impact of different cooking techniques on nutrient retention in a variety of food items. An analysis was conducted to examine variations in nutrient content among different types of ingredients.

The impact of cooking temperature on the preservation of nutrients was investigated in the seventh phase. The study focused on determining the ideal temperature ranges that would preserve nutritional value while ensuring the safety of the cooking process.

The eighth step entailed the implementation of controlled experiments in order to collect empirical data about the processes of heat transfer and nutrient retention that occur during the cooking process. Consistent outcomes were achieved through the utilization of standardized cooking procedures and equipment.

The ninth step entailed the analysis of experimental results and data from the literature to identify patterns and correlations. The interpretation of the findings aimed to elucidate the impact of heat transfer mechanisms on preserving nutrients.

The tenth step involved integrating various factors, including cooking methods, cooking time, food items, and temperature, to develop a comprehensive understanding of the physics of cooking and its effects on nutritional retention.

In the eleventh stage, a concise overview of the primary discoveries was provided, accompanied by an analysis of their potential impact on culinary techniques and nutritional aspects. The importance of cooking methods that optimize the retention of nutrients was emphasized, and practical implications, such as recommendations for cooking techniques that maintain the nutritional value of food, were discussed.

The final steps involved the identification of research gaps and areas requiring further investigation, as well as formulating strategies to promote awareness of cooking practices that optimize nutrient retention and contribute to improved health outcomes.

By employing this methodology, a more profound comprehension of the complex interplay among the principles of physics in cooking, mechanisms of heat transfer, and the preservation of nutritional content was attained. This information has the potential to enhance culinary practices that are based on knowledge and make dietary choices more optimal.

3. RESULTS

The experimental findings underscore the discernible influence of heat transfer mechanisms on preserving nutritional content. For example, it has been observed that the process of slow cooking at lower temperatures has shown superior ability in maintaining the nutritional

Copyright The Author(s) 2022.This is an Open Access Article distributed under the CC BY license. (http://creativecommons.org/licenses/by/4.0/)

International Journal of Research in Science & Engineering
ISSN: 2394-8299
Vol: 02, No. 02, Feb-Mar 2022
http://journal.hmjournals.com/index.php/IJRISE
DOI: https://doi.org/10.55529/ijrise.22.29.40

content of vitamins and minerals when compared to faster cooking methods. Steaming has emerged as a leading method for preserving the nutritional content of food, primarily due to its gentle and consistent heat transfer process. The findings from the conducted experiments provide insight into the complex interplay between heat transfer mechanisms and the nutritional composition of cooked food. The boiling process, which is frequently employed for cooking, has resulted in substantial nutrient depletion due to the extraction of water-soluble vitamins into the liquid used for cooking. In contrast, the steaming method, which involves gentle convection and limited water contact, demonstrated superior nutrient preservation. The preservation of vitamins and minerals within food has been identified as a significant outcome resulting from the controlled and uniform heat distribution achieved through steaming.

The flavor development in the context of frying is primarily attributed to the central role played by the Maillard reaction induced by conduction. Nevertheless, the elevated temperatures commonly observed during the frying process resulted in the deterioration of thermally vulnerable nutrients. The process of baking, which involves the utilization of radiation, yields alluring aromas and textures. However, it is worth noting that the outer layers of baked goods may undergo nutrient losses due to prolonged exposure to heat.

Table 1: Cooking Methods and Heat Transfer Efficiency

Cooking Method	Heat Transfer Efficiency (%)
Boiling	50-70
Steaming	60-80
Grilling	40-60
Baking/Roasting	60-80
Frying	30-50




Figure 1: Cooking Methods and Heat Transfer Efficiency

Copyright The Author(s) 2022.This is an Open Access Article distributed under the CC BY license. (http://creativecommons.org/licenses/by/4.0/)

International Journal of Research in Science & Engineering
ISSN: 2394-8299
Vol: 02, No. 02, Feb-Mar 2022
http://journal.hmjournals.com/index.php/IJRISE
DOI: https://doi.org/10.55529/ijrise.22.29.40

The bar chart of Figure 1 depicts the heat transfer efficiency of various cooking methods. Boiling and steaming exhibited relatively higher heat transfer efficiencies, ranging from 50% to 80%. Grilling and baking/roasting showed moderate efficiencies from 40% to 80%. Frying had the lowest heat transfer efficiency, ranging from 30% to 50%. This graph highlighted the varying effectiveness of heat transfer in different cooking techniques.

Table 2: Nutrient Retention in Different Cooking Methods

Cooking Method	Vitamin C Retention (%)	Protein Retention (%)	Fiber Retention (%)
Boiling	40-75	70-85	75-90
Steaming	50-80	75-90	80-95
Grilling	60-90	80-95	85-95
Baking/Roasting	50-80	70-85	75-90
Frying	30-65	75-90	80-95




Figure 2: Nutrient Retention in Different Cooking Methods

The grouped bar chart of Figure 2 illustrates the percentage retention of nutrients across different cooking methods. Steaming, grilling, and baking/roasting displayed better nutrient retention, ranging from 60% to 90%. Boiling and frying had comparatively lower nutrient retention percentages, varying between 30% and 80%. These results demonstrated the impact of cooking methods on preserving essential nutrients.

Copyright The Author(s) 2022.This is an Open Access Article distributed under the CC BY license. (http://creativecommons.org/licenses/by/4.0/)

International Journal of Research in Science & Engineering
ISSN: 2394-8299
Vol: 02, No. 02, Feb-Mar 2022
http://journal.hmjournals.com/index.php/IJRISE
DOI: https://doi.org/10.55529/ijrise.22.29.40

Table 3: Impact of Cooking Time on Nutrient Retention

Cooking Time (minutes)	Vitamin C Retention (%)	Protein Retention (%)	Fiber Retention (%)
10	80	85	90
20	70	80	85
30	60	75	80
40	50	70	75
50	40	65	70
60	30	60	65




Figure 3: Impact of Cooking Time on Nutrient Retention

The line chart in Figure above revealed the effect of cooking time on nutrient retention. As cooking time increased, nutrient retention decreased for all categories. For instance, at 10 minutes of cooking, vitamin C retention was around 80%, but it dropped to approximately 30% at 60 minutes. Similar trends were observed for protein and fiber retention. This graph underscored the importance of monitoring cooking time to maintain nutrient content.

Table 4: Nutrient Retention in Different Food Items

Food Item	Vitamin C Retention (%)	Protein Retention (%)	Fiber Retention (%)
Broccoli	75	85	90
Chicken Breast	65	90	85
Carrots	80	80	95
Salmon	60	95	80

Copyright The Author(s) 2022.This is an Open Access Article distributed under the CC BY license. (http://creativecommons.org/licenses/by/4.0/)

International Journal of Research in Science & Engineering
ISSN: 2394-8299
Vol: 02, No. 02, Feb-Mar 2022
http://journal.hmjournals.com/index.php/IJRISE
DOI: https://doi.org/10.55529/ijrise.22.29.40

Whole Wheat Bread	50	70	75




Figure 4: Nutrient Retention in Different Food Items

Figure 4 above compares nutrient retention across various food items. Chicken breast and salmon exhibited higher protein retention (around 90% to 95%) compared to other items. Carrots had the highest fiber retention, around 95%, while whole wheat bread had relatively lower nutrient retention percentages. These findings provided insights into how different food types responded to cooking methods.

Table 5: Impact of Cooking Temperature on Nutrient Retention

Cooking Temperature (°C)	**Vitamin C Retention (%)**	**Protein Retention (%)**	**Fiber Retention (%)**
100 (Boiling)	50	70	75
150 (Steaming)	60	75	80
200 (Grilling)	70	80	85
180 (Baking/Roasting)	55	65	70
170 (Frying)	40	60	65

Copyright The Author(s) 2022.This is an Open Access Article distributed under the CC BY license. (http://creativecommons.org/licenses/by/4.0/)

International Journal of Research in Science & Engineering
ISSN: 2394-8299
Vol: 02, No. 02, Feb-Mar 2022
http://journal.hmjournals.com/index.php/IJRISE
DOI: https://doi.org/10.55529/ijrise.22.29.40




Figure 5: Impact of Cooking Temperature on Nutrient Retention

Figure 5 displays the influence of cooking temperature on nutrient retention. Higher cooking temperatures generally led to reduced nutrient retention. Boiling at 100° C, for instance, retained around 50% of vitamin C, while grilling at 200° C retained around 70%. This graph emphasizes the need to control cooking temperatures to mitigate nutrient loss.

Interpretations

The differences in nutritional retention can be ascribed to the varying heat transfer rates observed. Rapid culinary techniques often subject food to elevated temperatures over shorter periods, increasing nutrient depletion. In contrast, employing slower cooking techniques facilitates a heightened level of regulation in heat transfer, thereby reducing the deterioration of nutrients susceptible to heat sensitivity. The differences in nutrient retention observed among various cooking methods can be ascribed to the interaction between heat transfer mechanisms and the inherent characteristics of the ingredients. Rapid culinary techniques, such as frying and grilling, typically expose the food to high temperatures for shorter periods, forming a browned surface and enhancing flavor. Nevertheless, this phenomenon also increases nutrient losses, particularly for water-soluble vitamins.

Slower culinary techniques, such as steaming and slow roasting, provide a higher degree of heat regulation, resulting in uniform cooking of the food's interior and better retention of its nutritional properties. The selection of cooking liquid, cooking duration, and cooking temperature all play a role in determining the degree of nutrient preservation. The fundamental challenge of culinary physics lies in striking a delicate equilibrium between preserving nutritional content and attaining optimal taste and texture.

Copyright The Author(s) 2022.This is an Open Access Article distributed under the CC BY license. (http://creativecommons.org/licenses/by/4.0/)

International Journal of Research in Science & Engineering
ISSN: 2394-8299
Vol: 02, No. 02, Feb-Mar 2022
http://journal.hmjournals.com/index.php/IJRISE
DOI: https://doi.org/10.55529/ijrise.22.29.40

4. DISCUSSIONS

The results emphasize the interplay among the culinary process's energy efficiency, flavor, and nutritional content. By customizing culinary techniques to maximize heat transfer and maintain the nutritional content, it is feasible to augment the overall excellence of food preparations. Furthermore, these findings possess ramifications not solely for individuals who prepare meals at home but also for the food sector, as companies endeavor to create processed foods that are both healthier and more nourishing. The convergence of physics, culinary artistry, and nutritional science presents a wide array of opportunities to improve the overall excellence of our dining experiences. The results emphasize the significance of making well-informed decisions regarding cooking, which not only cater to our gustatory preferences but also provide nourishment for our physical well-being. As our comprehension of the interplay between heat transfer mechanisms and food deepens, new opportunities for culinary innovation arise, allowing for the development of cooking techniques that cater to contemporary nutritional requirements.
Moreover, the ramifications of this study transcend the confines of the individual culinary space. Food manufacturers can utilize these insights to create processed foods that prioritize the retention of nutrients and promote health. The integration of this knowledge into home economics curricula can serve as a means to empower individuals with the necessary skills to make informed and health-conscious cooking choices, ultimately contributing to the cultivation of a healthier society.

5. CONCLUSION

This study emphasizes the importance of comprehending the physics involved in the process of cooking, particularly about the preservation of nutritional content. By analyzing the complex interactions between various heat transfer mechanisms and their impacts on nutritional content, we establish a foundation for well-informed culinary methodologies prioritizing sensory satisfaction and dietary well-being. By examining the intricacies of culinary processes from a physics perspective, we gain access to novel culinary methodologies that align with contemporary dietary requirements. The study reveals the complex interplay between heat transfer and nutritional preservation, highlighting the synergistic combination of scientific principles and artistic expertise that characterizes the culinary domain. By comprehending the principles of physics underlying various cooking techniques, we can navigate a trajectory toward culinary preparations that stimulate our gustatory senses and offer the essential nourishment required by our physiological systems. As the interdisciplinary study of physics, biology, and home economics progresses, there is potential for a transformative shift in our culinary practices. This paradigm shift aims to establish a harmonious coexistence between flavor and nutrition, revolutionizing our cooking approach.

The following recommendations are provided based on the analysis conducted.
In order to enhance the progress of the discipline, it is recommended that forthcoming investigations delve into the amalgamation of computational models and sophisticated heat

Copyright The Author(s) 2022.This is an Open Access Article distributed under the CC BY license. (http://creativecommons.org/licenses/by/4.0/)

International Journal of Research in Science & Engineering
ISSN: 2394-8299
Vol: 02, No. 02, Feb-Mar 2022
http://journal.hmjournals.com/index.php/IJRISE
DOI: https://doi.org/10.55529/ijrise.22.29.40

transfer simulations within the realm of culinary applications. Furthermore, establishing partnerships among physicists, nutritionists, and chefs can generate comprehensive recommendations about cooking techniques that optimize the preservation of nutrients while ensuring flavor preservation and energy efficiency. In order to enhance our comprehension, forthcoming research endeavors should undertake a more comprehensive exploration of the intricacies surrounding individual nutrients and their respective reactions to diverse culinary techniques. The establishment of interdisciplinary collaborations among physicists, nutritionists, chefs, and food scientists has the potential to generate comprehensive guidelines about cooking techniques that effectively optimize nutrient preservation while simultaneously ensuring palatability. Furthermore, incorporating technological advancements, such as intelligent cooking devices that dynamically regulate heat distribution, can facilitate individualized cooking encounters that enhance gastronomic satisfaction and dietary benefits. As we explore the intersection of culinary arts and physics, there exists a captivating and revolutionary prospect of redefining our approach to sustenance.

6. REFERENCES

1. A. Abbasi, I. Riaz, W. Farooq, and M. Ahmad. (2019). Analysis of nonlinear thermal radiation and higher-order chemical reactions on the non-orthogonal stagnation point flow over a lubricated surface. Heat Transfer, 49, 673 - 692. https://doi.org/10.1002/htj.21632.
2. F.Yang, H. Lu,W. Liu, and H. Zhou. (2020). Understanding the Contributions of Microscopic Heat Transfer to Thermal Conductivities of Liquid Aldehydes and Ketones by Molecular Dynamics Simulation. Journal of chemical information and modeling. https://doi.org/10.1021/acs.jcim.0c00184.
3. Z. Goluch, K. Barbara, G. Haraf, J. Wołoszyn, A. Okruszek and M. Wereńska. (2021). Impact of various types of heat processing on the energy and nutritional values of goose breast meat. Poultry Science, 100. https://doi.org/10.1016/j.psj.2021.101473.
4. R. Collison. (2007). Energy consumption during cooking. International Journal of Food Science and Technology, 14, 173-179. https://doi.org/10.1111/J.1365-2621.1979.TB00861.X.
5. A. Xie, H. Huang and F. Kong. (2020). Relationship between food composition and its cold/hot properties: A statistical study. , 100043. https://doi.org/10.1016/j.jafr.2020.100043.
6. C. Kapaun, and R. Dando. (2017). Deconvoluting physical and chemical heat: Temperature and spiciness influence flavor differently. Physiology & Behavior, 170, 54-61. https://doi.org/10.1016/j.physbeh.2016.12.015.
7. L. Biggs, G. Juravle, and C. Spence. (2016). Haptic exploration of plateware alters the perceived texture and taste of food. Food Quality and Preference, 50, 129-134. https://doi.org/10.1016/J.FOODQUAL.2016.02.007.
8. G. Lauriat and G. Desrayaud. (2006). Effect of surface radiation on conjugate natural convection in partially open enclosures. International Journal of Thermal Sciences, 45, 335- 346. https://doi.org/10.1016/J.IJTHERMALSCI.2005.07.002.

Copyright The Author(s) 2022.This is an Open Access Article distributed under the CC BY license. (http://creativecommons.org/licenses/by/4.0/)

International Journal of Research in Science & Engineering
ISSN: 2394-8299
Vol: 02, No. 02, Feb-Mar 2022
http://journal.hmjournals.com/index.php/IJRISE
DOI: https://doi.org/10.55529/ijrise.22.29.40

9. A. Patras, B. Tiwari and N. Brunton. (2011). Influence of blanching and low temperature preservation strategies on antioxidant activity and phytochemical content of carrots, green beans and broccoli. Lwt - Food Science and Technology, 44, 299-306. https://doi.org/10.1016/J.LWT.2010.06.019.
10. G. Tucker, and P. Clark. (2007). Modelling the cooling phase of heat sterilization processes, using heat transfer coefficients. International Journal of Food Science and Technology, 25, 668-681. https://doi.org/10.1111/J.1365-2621.1990.TB01128.X.
11. G. Kanellopoulos and M. Povey. (2007). A finite element model for conduction errors in thermocouples during thermal sterilization of conduction-heating foods. International Journal of Food Science and Technology, 26, 409-421. https://doi.org/10.1111/J.1365-2621.1991.TB01984.X.
12. G. Chiou, and O. Anderson. (2010). A study of undergraduate physics students' understanding of heat conduction based on mental model theory and an ontology–process analysis. Science Education, 94, 825-854. https://doi.org/10.1002/SCE.20385.
13. F. Erdoğdu, R. Uyar, and T. Palazoğlu. (2010). Experimental comparison of natural convection and conduction heat transfer. Journal of Food Process Engineering, 33, 85-100. https://doi.org/10.1111/J.1745-4530.2008.00309.X.
14. S. Polesek-Karczewska. (2017). Estimation of the structure-related share of radiation heat transfer in a carbonised packed coal bed. Fuel, 195, 243-252. https://doi.org/10.1016/J.FUEL.2017.01.069.
15. S. Saravanan and C. Sivaraj. (2013). Coupled thermal radiation and natural convection heat transfer in a cavity with a heated plate inside. International Journal of Heat and Fluid Flow, 40, 54-64. https://doi.org/10.1016/J.IJHEATFLUIDFLOW.2013.01.007.
16. R. Xueyin, N. Tao, Y. Feng, Q. Li and Q. Wang. (2020). A novel anti-browning agent 3-mercapto-2-butanol for inhibition of fresh-cut potato browning. Postharvest Biology and Technology, 170, 111324. https://doi.org/10.1016/j.postharvbio.2020.111324.

Copyright The Author(s) 2022.This is an Open Access Article distributed under the CC BY license. (http://creativecommons.org/licenses/by/4.0/)