
About The Nootropic Properties of Donsumine

Y.R. Mirzaev¹, E.M. Ruzimov^{2*}, R.A. Botirov³, S.F. Aripova⁴, I. Zhalalov⁵

^{1,2*,3,4,5}*Institute of Chemistry of Plant Substances named after Academician S.Yu. Yunusov of the Academy of Sciences of the Republic of Uzbekistan.*

Corresponding Email: ^{2*}tolmas4th@mail.ru

Received: 24 April 2023

Accepted: 11 July 2023

Published: 25 August 2023

Abstract: *Previous studies of Donsumine showed that Donsumine at doses of 3, 5, 10 mg/kg enhances motor activity (MA), verticalization, and at a dose of 100 mg/kg reduces it by 60%. It was also established that Donsumine at doses of 3 and 30 mg/kg enhances the activation of MA and verticalization caused by phenamine, the increase in MA persists until the end of the experiment, and when canceled it is compared with control mice. When canceled, the activity of experimental and control mice becomes the same. It was also established that Donsumine in the "open field" experiments at doses of 1, 10 and 100 mg/kg enhances the search and research activity in the open field according to C. Hall [1]. These data, which are elements of nootropic action, were the basis for a more detailed study of nootropic action. It is known that $\alpha 1$ adreno and dopamine receptors are the main ones in the realization of the nootropic action of substances. Our studies of the nootropic action Donsumine showed that Donsumine accelerated the production of conditioned reflex of passive avoidance (CRPA), the development of the esophageal reflex in the classical maze, improved spatial memory in the water maze according to Morris. He showed antagonism to the amnestic action of scopolamine, the development of the esophageal reflex, as well as in experiments in the water maze according to R. Morris [2]. The antagonism of Donsumine and scopolamine was competitive.*

Keywords: *Motor Activity, "Open Field", Scopolamine, CRPA, Amnestic Action, Research Activity.*

1. INTRODUCTION

Arundo (reed) or *Arundo donax* L. for many years it has been used not only for agricultural or household needs, but also for preparation of tincture from its leaves, is used in oriental folk medicine as a diuretic, for the treatment of sexual diseases, as well as for diseases such as ascites, candidiasis, stomatitis, decoctions of dried crushed parts. This type of plant is much more often cultivated in countries such as Romania, Italy, France, it is widely distributed in the Bukhara, Kashkadarya and Ferghana regions of our country scattered. In our country, as



well as all over the world, academicians A.P. Orekhov and S.Yu Yunusov's have been for many years continuation of scientific research, Arundo Donax L. A number of scientific research works are carried out on the basis of the plant [13-18]

The forage plant reed (Arundo donax L) is a source of indole alkaloids of the tryptamine series [3] and, first of all, donaxin, in which selectivity to 5HT_{2A} receptors of the serotonin receptor subtype has been revealed [4]. According to T. Mestre [5], 5HT_{2A} are a source of psychotropic drugs, including nootropic ones. In particular, at the Institute of Chemistry of Plant Substances, academician named after S.Yu. Yunusov of the Academy of Sciences of Uzbekistan Arundo Donax L. in continuation of a series of research works based on the plant for many years, the nootropic effect of the amount of alkaloids on the surface of this plant was studied [13]. Thus, the comparison of anticonvulsant, antidepressant and anxiolytic activity of recently synthesized tabium and synthetic structured substances with some drugs that are widely used in practice under experimental conditions is being studied. As a direct continuation of this research work, screening studies study the biological activity of substances isolated from plants, with their large-scale distribution in our country and Asia [19, 20]

In this regard, our studies of Donsumin, in addition to the psychostimulating effect, also revealed the anxiolytic, as well as the described nootropic effect.

2. PHARMACOLOGICAL RESEARCH METHODS AND MATERIALS

All studies to determine nootropic activity were conducted on mongrel laboratory white mice with a body weight of 20-22 g and on mongrel laboratory white rats with a body weight of 180-220 g, which were cared for and kept in standard quarantine conditions for 14 days [21].

1.1. The effect of Donsumine on D₂ of the stimulating effect of phenamine. It is known that phenamine stimulates D₂ dopamine receptors, which are involved in the process of cognitive functions of animals and humans. [6, 7] is mediated with the participation of DA₂ α_1 adreno receptors in the form of verticalization (strut) and enhancement of MA. In this experiment, the effect of Donsumine on the sensitivity of DA₂ receptors was studied. The experiments were carried out on 3 groups of mice using the I. P. Lapin technique [8],

The experience consisted of: the number of verticalization and intersection of the lines of MA mice in the control was calculated. against the background of D₂ stimulating effect of phenamine from a dose of 5 mg / kg n / a and the same experience on the background of Donsumine administered 2 hours before administration of phenamine. At the same time, the effect of the drug on the severity of verticalization was determined.

1.2. Study of search and research activities in the open field by Hall. The mice landed on an open area divided into 5x5 squares and drilled holes D = 15 mm. The number of contacts of mice with squares was estimated as the search activity of mice, and the contacts of mice with minks (peeping) were taken as research activity.

1.3. The effect of Donsumine on the development of the conditioned reflex of passive avoidance (CRPA) was evaluated in a 2-chamber block, of which one chamber was light and had an entrance to a dark chamber with an electrode floor, which was connected to the electrical stimulator ESU 01-1. In the experiment, mice landed one animal at a time in a light compartment with their tail pointing towards a dark compartment, from where they, as a rule,



moved into a dark compartment with an electrode floor. After 5 seconds, an alternating electric discharge (50 volts, 0.5 mA, 20 Hz) was applied to the floor electrodes.

1.4. The influence of Donsumine on the speed of the conditional motor reflex was determined in a classic maze with 2 chambers. The experiments were carried out for 3 days. In the evening, the rats were deprived of food with access to water. In the morning, an experiment began on finding a compartment with food by rats, sweetened milk with bread in a 100 ml Petri dish. On the first day, the rats were trained three times to find food, while the time of finding the final compartment with food was recorded.

1.5. The effect of (E) on the spatial memory of mice was studied in a water maze using the Morris method. The water maze consisted of a galvanized tank D = 40cm and 30 cm high. At the opposite end of the tank there is a wooden platform D = 5 cm, immersed in water with a temperature of 27-29 C at 0.5-0.7 cm below the water surface. To ensure the opacity of the water, milk with a volume of 300 ml of 3.2% fat content was poured.

The time of the platform 's location was recorded

1.6. Effect of Donsumine on the amnesia of mice caused by the administration of scopolamine 1 mg/kg s/c.

The discussion of the results obtained on the basis of the conducted studies was carried out in comparison with the control group and statistical processing of the results was performed using the methods presented in R.B.Strelkov [22].

3. THE RESULTS OF THE STUDY

1.1. The effect of Donsumin on MA. Previous studies have shown that Donsumine starting with a dose of 1 mg/ kg, the behavior of verticalization also increases. At a dose of 3 mg/kg, these effects increased 2-3 times. At a dose of 10 mg/kg, they were additionally increased by 45%, i.e. relatively less than from a dose of 3 mg/kg. It must be assumed that Donsumine in these doses enhanced MA and verticalization due to the activation of DA2 dopamine a1 receptors [6, 7,] which participate in the manifestation of the nootropic effect of Ginko Biloba, Piracetam and others.[9, 10, 11, 12, 13]

1.2. The above data were confirmed in experiments on the influence of Donsumine on search and research activity in experiments in an open field by Hall. The following studies are designed to confirm the participation of Donsumine nootropic action.

1.3. The effect of donsumine on the development of the conditioned reflex of passive avoidance of punishment (CRPA). The experiments were carried out on 2 groups of mice with 6 animals in a 2-chamber block with dark and light compartments. It was found that in the control group of mice, when re-placed in a light chamber, out of 6 mice, only 2 did not enter the dark chamber, i.e. they developed an avoidance reflex.(33%) . In the second experimental group of mice that were previously injected with Donsumine at a dose of 5 mg/kg 24 and 2 hours before the start of the experiment, none entered, i.e. the avoidance reflex was developed in all mice.

1.4. The effect of Donsumine on the latent period of the food-motor reflex in experiments on rats in a maze. On the first day of the experiment, each rat was trained to run through the maze 3 times. The next day, each rat was passed through the maze and the duration of the run was used as control data.



The 2nd and 3rd runs were carried out 2 and 4 hours after administration of Donsumine 3 and 10 mg/kg. It was found that the latency period in three groups of rats averaged 15, 17, 18 seconds. After 2 hours, the runs were repeated, and the latent periods were from a dose of 3 mg/kg-8 seconds, from a dose of 10 mg/kg-10 seconds, and in controls-14 seconds.

The results of the experiments indicate that the duration of the run in the control group did not change much, from a dose of 3 mg/kg, the latency period decreased from 17 to 8% or by 53%; from a dose of 10 mg/kg, it changed from 18 seconds to 44%.

The experiments confirmed the nootropic effect of Donsumine was confirmed, and the dose of 3 mg/kg was the most active.

1.5. The effect of Donsumine on the amnesic effect of scopolamine. On the third day, the experiment was repeated, but the second group after Donsumine 3 mg/kg after 1 hour was injected with scopolamine 1 mg/kg s/c., the third group was still 10 mg/kg s/c. As a result, in the 1st group of mice, the latency period decreased from 15 seconds to 14 seconds, i.e. practically unchanged; in the second group, against the background of Donsumine 3 mg/kg + scopolamine 1 mg/kg, there was a less pronounced shortening of the latency period. i.e., in the second group, the latency period after scopolamine and Donsumine in the indicated the dose was from 18sec, i.e. decreased to 14 seconds, i.e. the latency period was longer than in the second experiment, which proved the mutual antagonism of Donsumine and scopolamine. In group 3, the latency period from Donsumine 10 mg/kg remained the same, i.e. did not change, was the same as in the previous experiment.[2.4] The data provided indicate that the latency period has remained the same.

The results of the experiment indicate the competitive antagonism of scopolamine and Donsumine.

1.6. The effect of Donsumine on the latent period of the platform in the water maze in Morris experiments on mice.

In these experiments, the effect of substances on spatial memory is studied. It was found that the average latency period of the platform in the control 6 mice was 126 seconds. In the groups of mice with the introduction of Donsumine 3 and 10 mg/kg inside, the time of finding the maze was 56, 65 seconds, i.e. about 2 times faster.

In experiments with a water maze, in a group of mice with the introduction of Donsumine 3 mg /kg, scopolamine 1 mg/kg was additionally injected, and in a group with the introduction of Donsumine 10 mg/kg, distilled water 0.2 ml/kg was injected. The speed of finding the platform in control mice was 114 seconds, against the background of Donsumine + scopolamine, the time of finding the platform was 86 seconds, and in the 3rd group against the background of Donsumine 10 mg/kg was 110 seconds. Thus, in this experiment, the 2nd group of mice in which Donsumine and scopolamine were injected extended the time of finding the maze from 56% to 114%, thus scopolamine and Donsumine counteracted each other, i.e. showed mutual antagonism. In group 3, the shortening of the latency period was almost no different from the results of the previous experiment.

From the data presented, it can be seen that scopolamine lengthened the latency period by 2.5 times, and the preliminary administration of Donsumine the latency period approached the control level. It can be seen from the experimental data that Donsumine has an anti-amnesic or nootropic effect.



4. DISCUSSION

Thus Donsumine, which is a fraction of indole alkaloids of the tryptamine series isolated from reed reed, is a new original nootropic agent whose mechanism is the activation of α_1 adreno and D2 dopamine receptors plus, presumably, the blockade of 5HT_{2A} serotonin receptors. Nootropic effect manifested in a test to accelerate the production of CRPA in mice, to accelerate the food-motor reflex in a classic maze, in a test for finding a water maze by Morris. The nootropic effect of Donsumine, was antagonized by the amnesic effect of scopolamine in the last two tests. The nootropic effect of Donsumine was manifested in doses of 3-10 mg/kg of 1/300 – 1/100 of LD₅₀. Doses up to 100 mg/kg orally also had a nootropic effect, but it was combined with muscle relaxation from large doses of Donsumine.-100 mg/kg. Donsumine is a representative of a new chemical class of new nootropic drugs.

5. CONCLUSIONS

1. Donsumine is a fundamentally new nootropic agent.
2. Fractions of indole alkaloids of the tryptamine series are Donsumine, consisting of Donaxin (60-80%), Ardine, Donaxamine, Arundine. Donsumine consisting of these alkaloids has a nootropic effect.
3. The mechanism of nootropic action of Donsumine involves α_1 adreno- DA2 dopamine receptors involved in the mechanism of action of Ginko, Piracetam.
4. Donsumine and scopolamine showed antagonism to each other's effects by competitive type.

6. REFERENCES

1. C. Hall, J. Comp. Psychol., 22, 345-352 (1936).
2. Morris R. Developments of a water-maze procedure for studying spatial learning in the rat. J Neurosci Methods. 1984, V. 11, P. 47–60.]
3. З.В.У.Хужаев, Автореф.дис.д-рахим.наук,Ташкент(2005).
4. Frolidi G, Silvestrin B., Dorigo P., Caparrotta L. Gramine: a vasorelaxing alkaloid acting on 5-HT_{2A} receptors. //Planta Med.2004 Apr; 70(4):373-375.
5. T. Mestre, M. Zurowski, S. Fox, Expert Opin. Investig. Expert Opin Investig Drugs, 22 (4):411-21 (2013). doi: 10.1517/13543784.2013.769957. Epub 2013 Feb.
6. B. Rothman, M. Baumann, Amphetamine-type central nervous system stimulants release norepinephrine more potently than they release dopamine and serotonin†
7. Protais P., Costentin J., Schwartz J.C.// Climbing behavior induced by apomorphine in mice: a simple test for the study of dopamine receptors in striatum. Psychopharmacology (Berl.) 1976 Oct 20; 50(1):1-6.
8. Лапин И.П., Слепокуров М.В. Анксиогенная активность фенилэтиламина в тесте социальной изоляции на мышах //Фармакол. и токсикол., 1991, т. 54, № 6, с. 9-11.
9. K.D. Alex and E.A. Pehek1, Pharmacologic mechanisms of serotonergic regulation of dopamine neurotransmission Pharmacol Ther. 2007 Feb; 113(2): 296–320.



10. B Søholm/Clinical improvement of memory and other cognitive functions by Ginkgo biloba: review of relevant literature//Adv Ther 1998 Jan-Feb;15(1):54-65.
11. Bruce J Diamond 1, Mary R Bailey/Ginkgo biloba: indications, mechanisms, and safety//Psychiatr Clin North Am 2013 Mar;36(1):73-83.
12. Guoyan Yang, Yuyi Wang, Jin Sun, Kang Zhang, Jianping Liu 1/Ginkgo Biloba for Mild Cognitive Impairment and Alzheimer's Disease: A Systematic Review and Meta-Analysis of Randomized Controlled Trials//Curr Top Med Chem 2016;16(5):520-8
13. Мирзаев Ю.Р., Рузимов Э.М., Арипова С.Ф., Жалолов И.Ж. (2022). Скрининг тадқиқотларда алкалоидлар йиғиндиси "донсумин" нинг ноотроп таъсирини ўрганиш. Eurasian journal of medical and natural sciences, 2(11), 256–262. <https://doi.org/10.5281/zenodo.7221072>
14. Jalolov I.J., Khujaev V.U., Aripova S.F., Mirzaev Yu.R., Kurmikov A.G. / Chemistry, transformations and. Pharmacological activity of Arundo donax L. alkaloids. // Nitrogen- containing heterocycles and alkaloids. -(Iridium Press). - Moscow. -2001. –V.1. -P. 311-313.
15. Арипова С.Ф., Хужаев В.У., Жалолов И.Ж., Сагдуллаев Ш.Ш. / Алкалоиды гигантского злака Arundo donax L. химия, структура, свойства, технология // Монография. Ташкент. -2017. С. 256.
16. Ботиров Р.А., Саноев З.И. Математическое планирование процесса экстракции алкалоида донаксина из растения Arundo donax L // Universum: Химия и биология : электрон. научн. журн. 2018. № 7(49). URL: <http://7universum.com/ru/nature/archive/item/6065>
17. Мирзаев Ю.Р., Арипова С.Ф. Арундо-донакс - как потенциальный источник серотонинэргических соединения. 5-й Международный симпозиум химии природных соединений. Ташкент, 2003. С. 65.
18. Мирзаев Ю.Р., Арипова С.Ф. Сравнительное фармакологическое исследование серотонина и донаксина. 2-й Международный симпозиум по съедобным растительным ресурсам и биоактивным ингредиентам. Урумчи, Китай, 2010. С. 122-123.
19. Т.Т. Ҳамроев, Н.М. Маматқулова, З.И. Саноев, С.З. Рашидов, И.Т. Абдиназаров, П.А. Нурмахмадова, Н.Қ. Хидирова, У.М. Якубов. (2022). Adonis turkestanica ўсимлигининг экстракция жараёнида ҳосил бўлган қолдиқ моддаларнинг анксиолитик фаоллигини скрининг тадқиқотларда ўрганиш. Eurasian journal of medical and natural sciences, 2(12), 146–152. <https://doi.org/10.5281/zenodo.7332882>
20. Т.Т.Ҳамроев, Н.М.Маматқулова, П.А.Нурмахмадова, С.З.Рашидов, И.Т.Абдиназаров, С.Д.Раҳимбоев, Н.Қ.Хидирова, У.М.Якубов. (2022). Adonis turkestanica ўсимлигининг экстракция жараёнида ҳосил бўлган қолдиқ моддаларнинг ўткир захарлилиги ва биологик фаоллигини скрининг тадқиқотларда ўрганиш. Eurasian journal of academic research, 2(12), 447–454. <https://doi.org/10.5281/zenodo.7332870>



21. National Research Council. 2011. Guide for the Care and Use of Laboratory Animals: Eighth Edition. Washington, DC: The National Academies Press. <https://doi.org/10.17226/12910>.
22. Стрелков Р.Б. Статистические таблицы для ускоренной количественной оценки фармакологического эффекта. Фармакология и токсикология 1986. №4 с.100-104.