



Creating a Score to Assess Weighting Impacts of Smartphone Addiction among Health and Medical Students

Ammar Ali Asghar Khalil^{1*}, Hewa Sattar Salih², Abdulkhaleq A Ali Ghalib³

¹Kirkuk Health Directorate, Kirkuk City, Iraq.

²Professor (Dr.) Hewa Sattar Salih; College of Nursing, University of Kirkuk, Kirkuk city, Iraq.

³Emeritus Professor (Dr.) Abdulkhaleq A Ali Ghalib, College of Health and Medical Technology, Baghdad Iraq.

Email: ²Dr.hewa-salih@uokirkuk.edu.iq, ³abdulkhaliq.alnaqeeb@yahoo.com

Corresponding Email: ^{1*}ammr.charely20@gmail.com

Received: 19 May 2024

Accepted: 03 August 2024

Published: 18 September 2024

Abstract: *Aim of Study: This paper is aimed to introducing a new creating measurement score to estimating a significant of an extracted factors that reflects weighting of its components in terms of proposed studied domains for assessing status of impacts Smartphone addiction, in a way that achieving to the core of the evaluation process for new observation misuse of Smartphone among medical faculty students.*

Methods: *A cross-sectional design using a convenient sampling method which were selected of data, of sample size 772 participants from different studying stages concerning of medical and health faculties in Kirkuk University. Smartphone addiction scale (SAS), Nordic questionnaires for the analysis of musculoskeletal symptoms and the severity of Digital Vision Syndrome (DVS) was measured using a twelve-item was using. Data was analyzed using factor analysis of principle components method by using (SPSS) application, version 22. Results: The study sample included 347 males (44.9%) and 425 females (55.1%), with an average age of 21.09 ± 1.75 . Primary assessments showed highly significant relationships ($P < 0.01$) among the studied domains. The extracted factors revealed two dominant components, accounting for 67.22% and 32.78% of the total covariance, named the "Psycho-Socio Impact Factor" and "Physical Impact Factor," respectively. These findings offer a clear and robust understanding of the interrelation between the studied domains.*

Conclusions: *Factor analysis identified two main factors: the psycho-socio factor, related to family, environment, and psychological aspects, and the physical impact factor, linked to vision and musculoskeletal health. These findings suggest that interventions for smartphone misuse should target these interconnected factors to reduce negative effects. Additionally,*



the research highlights the importance of these domains in developing health-related quality of life programs, awareness workshops, and research into acquired autism in children.

Keywords: *Smartphone Addiction, Psychological Problems, Social Problems, Physical Problems.*

1. INTRODUCTION

In fact, the development of Smartphones over the past few years has made them more accessible and has led to an increase in their use by a diverse spectrum of people worldwide (Achangwa et al., 2022). Increasing reliance of society on Smartphones has led to the emergence of a new problem known as Smartphone addiction (Sarhan, 2024). Smartphone addiction is defined as the lack of ability to regulate Smartphone usage, even in the face of detrimental consequences for the users (Cha S & Seo, 2018). College students endure elevated levels of stress, potentially stemming from the frequency of examinations, extensive and laborious academic workload, and apprehension regarding the future (Gazzaz et al., 2018). When confronting with these stresses, University students are more prone to becoming addiction to Smartphone technologies. Smartphone usage for Internet access and excessive video game playing are linked to lower academic achievement (Mehammed-Ameen et al., 2018). It also effects on energy level, and food and exercise habits are all negatively impacted by Smartphone addiction. It also increases anxiety, sadness, insomnia, emotional instability, and even increases the risk of suicide due to cyberbullying and online fraud (Wang et al., 2023). Every university campus is linked to the Internet (I. Mohammed Ali & Nihad, 2021; Nife et al., 2022). Smartphones are commonly referred to as "handheld internet" devices (Lee et al., 2020). A person develops Smartphone addiction when they use their phone excessively, which can have detrimental repercussions on their life and their physical and emotional well-being (Shahrestanaki2020). Individuals who actively participate in social media and frequently use Smartphones are more prone to experiencing elevated levels of hopelessness and thoughts of suicide (Twenge et al., 2018). moreover, physical issues, such as musculoskeletal problems, migraine headaches, neck and wrist pain, unclear eyesight, pinch strength, and hand function (Kim, 2013; Kee et al., 2016). Addiction to Smartphones has four main characteristics: withdrawal, compulsion, functional impairment, and tolerance. Common symptoms of Smartphone addiction include anxiety, impaired function, and difficulty focusing and making decisions (Liu et al., 2022). University student marks a crucial phase where one starts to physically distance themselves from family and friends, leading to the realization of loneliness. Loneliness is a prevalent experience that can be found among Smartphone users. Individuals develop with psycho-social issues to social isolation and loneliness, due to their excessive use Smartphone (Karaoglan Yilmaz et al., 2023). Addiction to Smartphones has an impact on mental health Because people with depression or anxiety use Smartphones as a coping mechanism to deal with depressing and negative emotions, melancholy and anxiety can lead to technology addiction (Cha S & Seo, 2018) .as well as to the adverse psychological impacts, prolonged use of Smartphones can also lead to impaired vision (Sohn et al., 2019). Smartphone s emit electromagnetic radiation, including non-ionizing and ultraviolet UV radiations. Excessive use of Smartphones can increase the exposure of the eye to this radiation, leading



to biological damage to the eye cells or the appearance of defects. In addition, inadequate tear lubrication can lead to the development of dry eye disease, which can cause various eye conditions (Ismael, 2022). Recognizing the gravity of Smartphone addiction and its increasing prevalence, Nevertheless, there is a dearth of research on the prevalence of Smartphone addiction among university students in the country. Therefore, additional research on medical students is necessary to address the changes occurring in the healthcare and education system. This study aims to Creating new weighted score for assessing addiction status impact of new observation misuse of Smartphone among "Medical Faculty Students".

2. LITERATURE REVIEW

Smartphone addiction leads to disruptions in daily life, with increased device use over time and anxiety when separated from the phone (Lee & Cho, 2015). Smartphone addiction is believed to be derived from Internet addiction because of the similarity in symptoms and detrimental impact on users. Internet addiction is a condition that falls under the category of impulse control disorders. It is characterized by excessive and pathological usage of the Internet. Smartphone addiction can be classified as a behavioral addiction, similar to Internet addiction (Cha S & Seo, 2018). Research has demonstrated that excessive use of screen-based devices can lead to behavioral issues such as impulse control disorders, less social engagement, decreased physical activity, and difficulties in relationships (Basheer et al., 2021).

This may involve conflicts, diminished familial engagement, and emotional detachment. When a Smartphone is used excessively in the outside world, it can negatively impact social connections, work or school performance, and overall involvement with the external environment. Furthermore, the significant correlation with psychological concerns implies that Smartphone addiction is connected to mental health difficulties, including anxiety, despair, and stress. These findings suggest that Interventions aimed at reducing Smartphone addiction should focus on improving family dynamics, social interactions, and providing psychological support. The significance of the physical impact factor Excessive Smartphone use can result in notable vision issues, including eye strain, dryness, and potentially enduring harm. Additionally, the significant strain on the musculoskeletal system indicates physical problems such as neck discomfort, back pain, and other related conditions caused by improper posture and lengthy periods of use, it is consistent with the results of a study conducted in Malaysia in 2020 by Hassim et al. these physical health issues Efforts to mitigate Smartphone addiction should include strategies to alleviate vision and musculoskeletal problems, such as promoting proper ergonomics and regular breaks.

3. METHODS

A cross-sectional design using of convenient sampling method, which were selected of data at Feb. 2024, with sample size 772 participants from different studying stages concerning of medical and health faculties in Kirkuk University.

Four colleges were included: Medicine, Dentist, Pharmacy and Nursing college, with a sampling rate of 20% of the students sampling population had been selected for this study. The study excluded postgraduate students and those who did not agreed to be participate. A



proposed questionnaire was used, that consisted of three parts, the first was dedicated for the socio-demographical characteristics variables (SDCv.) of studied sample, and the second was dedicated for the risk factors associated with addiction misuse of Smartphone, and the third part was dedicated five main domains, such as: Family environment, External environment, Health problems related to vision, Health problems related to musculoskeletal pain, and Psychological, and that depending mainly on what was scored and published by (Abdulkhaleq A Ali Ghalib, et al., 2012).

Also, Smartphone addiction scale (SAS) after taking permission from author of the instrument (Kwon et al., 2013). Nordic questionnaires for the analysis of musculoskeletal symptoms (Kuorinka et al., 1987), and the severity of Digital Vision Syndrome (DVS) was measured using a twelve-item (Hundekari et al., 2021).

The approval was obtained from the students with an explanation of the research objectives and the importance of the study.

The English, Arabic and Kurdi paper-based version of proposed questionnaires was distributed to participants. Participants were assured that their information would be kept confidential and that there was no need to provide any identifying information. Data have been collected and the duration of each interview was between 10-20 min. The students answered the questionnaire in a convenient space in their classroom. using Likert scoring scales of 3 differentiated categories, such that: "Never, Sometimes, and Always". Reliability of the questionnaire showed that a very high level of stability and internal consistency of the studied subjective at the level of items of the applied questionnaire. In this study, Cronbach's α coefficient was (0.8853).

Statistical Analysis:

The statistical data analysis technique approaches were used to analyze and assess the study results using the statistical package for social science (SPSS) ver. (22.0) Descriptive data analysis included (Frequencies, Percentage, mean of score (MS), Standard Deviation (SD), Person's correlation coefficients, and inferential procedures were applied, such that: The Chi-Square test, Binomial test, The contingency coefficients (C.C.) test. Factor analysis of using principal components method. The P-value < 0.05 in at least were considered statistically significant.

4. RESULTS AND DISCUSSION

Table 1: Distribution of Socio-Demographic Characteristics Variables of studied sampling population with Comparison' Significant

SDCv.	Classes	No.	%	C.S. (*) P-value
Gender	Male	347	44.9	P=0.006 (HS)
	Female	425	55.1	
Age Groups Yrs.	< 20 yrs.	177	22.9	$\chi^2= 152.67$ P=0.000 (HS)
	20 21	213	27.6	
	22 23	342	44.3	
	> 24 yrs.	40	5.2	



	Mean ± SD	21.09 ± 1.75		
Marital Status	Married	30	3.9	P=0.000 (HS)
	Single	742	96.1	
Residency	Urban	719	93.1	P=0.000 (HS)
	Rural	53	6.9	
Socio-Economic Status	Low	95	12.3	$\chi^2= 152.67$ P=0.000 (HS)
	Moderate	502	65.0	
	High	175	22.7	
Rate of hours using smart phone per 24hrs?	1_3	65	8.40	$\chi^2= 261.11$ P=0.000 (HS)
	3_5	277	35.9	
	> 5	430	55.7	
Rate of sleeping hours during a day?	1_3	513	66.5	$\chi^2= 381.87$ P=0.000 (HS)
	3_5	119	15.4	
	> 5	140	18.1	
Body Mass Index BMI	Under weight	4	0.50	$\chi^2= 668.155$ P=0.000 (HS)
	Normal weight	407	52.7	
	Overweight	334	43.3	
	Obese	27	3.50	
	Total	772	100	

HS: Highly Sig. at P<0.01; Testing based on One-Sample Chi-Square test, and Binomial test.

The total number of participants in this study was 772, with 347 (44.9%) males and 425 (55.1%) females. The subjects who participated aged between 18 to 26 years, The average age was 21.09 ± 1.75. Regarding marital status, the majority of the analyzed sample were classified as "Single", accounting for 742 individuals (96.1%), while the remaining individuals were classified as married. the majority of the participants 719 (93.1%), were classified as having urban residency. moreover, socio-economic status almost three quarters of the sample into the "Low" and "Moderate" categories based on the WHO score, accounting for 597 individuals (77.3%), while leftover was assigned high level. 707(91.6%) of participants were used Smartphone at 3hrs and more. One-third of the sample (259 people, 33.5%) sleep 3 or more hours daily, while the remaining 513 (66.5%) sleep less than 3 hours. nearly half of the sample (361 people, 46.8%) are overweight or obese, with highly significant differences P<0.01 (Table1).

Table 2: Distribution of Favorite and Purposes of Smartphone Application used

Favorite Smart Phone Application		Response	No.	%	C.S. (*) P-value
Favorite Applications	Face book	None	457	59.2	P=0.002 (HS)
		Favorite	315	40.8	
	Snap chat	None	403	52.2	P=0.235 (NS)
		Favorite	369	47.8	
	Instagram	None	66	8.5	P=0.002 (HS)
		Favorite	706	91.4	



Purpose of Smart phone used	YouTube	None	324	42.0	P=0.002 (HS)	
		Favorite	448	58.0		
	Play games	None	456	59.1	P=0.002 (HS)	
		Favorite	316	40.9		
	Google	None	526	68.1	P=0.002 (HS)	
		Favorite	246	31.9		
	Twitter	None	605	78.4	P=0.002 (HS)	
		Favorite	167	21.6		
	Study	Study	None	140	18.1	P=0.002 (HS)
			Favorite	632	81.9	
		Communication	None	110	14.2	P=0.002 (HS)
			Favorite	662	85.8	
Entertainment		None	136	17.6	P=0.002 (HS)	
		Favorite	636	82.4		

Hs: Highly Sig. At P<0.01; Testing Based On Binomial Test.

The most favored smartphone applications, ranked from highest to lowest, are Instagram (91.4%), YouTube (58.0%), Snapchat (47.8%), gaming (40.9%), Facebook (40.8%), Google (31.9%), and Twitter (21.6%). The primary purposes for using smartphones, in order, are communication (85.8%), entertainment (82.4%), and study (81.9%). Significant differences were found between the observed and expected frequencies of smartphone use for all purposes (P<0.01).

Table (3): Summary Statistics of Student's responding items concerning of Social Problems (Family Environment) domain (N=772)

Main Domains	MS	SD	RS%
Family Environment			
I suffer from a lack of family support	1.58	0.70	52.67 L
My relationship with my family is not what it used to be	1.65	0.59	55.00 L
I do not participate with my family in social activities	1.81	0.65	60.33 M
I ignore my family members when I use a Smartphone	1.87	0.68	62.33 M
I experience a decrease in face-to-face contact with my family members	1.84	0.70	61.33 M
I feel comfortable staying alone at home	2.14	0.73	71.33 M
I became less interested in other daily tasks	2.06	0.67	68.67 M
I miss some important conversations and family events	1.97	0.68	65.67 M
Using Smartphone even in the main meals	1.91	0.71	63.67 M
I feel that my relationship with my family has changed	1.72	0.61	57.33 M
Preferring talking with my Smartphone buddies to hanging out with my real-life friends or with the other members of my family	1.72	0.60	57.33 M
(ExternalEnvironment)			



Missing planned work due to Smartphone use	2.10	0.73	70.00 M
Having a hard time concentrating in class, while doing assignments.	2.16	0.64	72.00 M
Having a hard time concentrating in working due to Smartphone use	2.06	0.68	68.67 M
I find it difficult to deal with others	1.73	0.62	57.67 M
I don't want to talk to others	1.7	0.64	56.67 M
I feel lonely and isolated from others	1.69	0.66	56.33 M
I feel like I don't have close friends	1.54	0.65	51.33 L
I feel unaccepted by others	1.55	0.69	51.67 L
I think using Smartphone negatively affects in my score in university	2.21	0.7	73.67 M
Preferring searching from my Smartphone to asking other people	1.93	0.61	64.33 M
Feeling that my Smartphone buddies understand me better than my real-life friends	1.75	0.68	58.33 M
Feeling that my relationships with my Smartphone buddies are more intimate than my relationships with my real-life friends	1.71	0.64	57.00 M
Health problems related to vision			
Experiencing light headedness due to excessive Smartphone use	1.80	0.71	60.00 M
Experiencing headache due to excessive Smartphone use	1.77	0.66	59.00 M
Experiencing blurred vision due to excessive Smartphone use	1.56	0.65	52.00 L
Experiencing double vision due to excessive Smartphone use	1.15	0.40	38.33 L
Tiredness in eyes	1.92	0.62	64.00 M
Straining of eyes	1.94	0.62	64.67 M
Heaviness in eyes	1.88	0.64	62.67 M
Burning sensation in eyes	1.76	0.66	58.67 M
Eye redness	1.65	0.65	55.00 L
Watery eyes	1.33	0.57	44.33 L
I feel ocular pain	1.83	0.72	61.00 M
Excessive use leads to weakness in the process of attention and concentration	1.97	0.60	65.67 M
Lack of attention to the radiant blue light from a Smartphone screen has the effect of a disruption of the biological clock in the body	1.93	0.57	64.33 M

Continue ...

Main Domains	MS	SD	RS%
--------------	----	----	-----



Health problems related to Musculoskeletal Pain			
Neck	2.12	0.70	70.67 M
Shoulder	1.94	0.68	64.67 M
Upper back pain	1.85	0.69	61.67 M
Elbow	1.61	0.67	53.67 L
Wrist/Hand	1.62	0.69	54.00 L
Lower back pain	1.56	0.63	52.00 L
Hips/Thigh	1.3	0.53	43.33 L
Knees	1.25	0.48	41.67 L
Ankles/Feet	1.28	0.53	42.67 L
Psychological Problems			
I feel like I don't have the energy to do my work	2.21	0.71	73.67 M
I feel uninterested in everything	2.00	0.61	66.67 M
I feel sad and depressed	1.87	0.63	62.33 M
I feel like I don't have any fun doing anything	1.71	0.66	57.00 M
I feel like life has no meaning	1.58	0.70	52.67 L
Feeling tired and lacking adequate sleep due to excessive Smartphone use	2.11	0.72	70.33 M
Being able to get rid of stress with a Smartphone	1.99	0.61	66.33 M
Having my Smartphone in my mind even when I am not using it	1.93	0.68	64.33 M
Feeling impatient and fretful when I am not holding my Smartphone	1.96	0.69	65.33 M
Getting irritated when bothered while using my Smartphone	1.96	0.71	65.33 M
Constantly checking my Smartphone so as not to miss conversations between other people on Twitter or Face book	2.25	0.73	75.00 M
Checking SNS (Social Networking Service) sites like Twitter or Face book right after waking up	2.40	0.71	80.00 H
Feeling the urge to use my Smartphone again right after I stopped using it	2.42	0.69	80.67 H
Having tried time and again to shorten my Smartphone use time, but failing all the time	2.01	0.77	67.00 M

RS%: Relative Sufficiency Assess By (L: Low; M: Moderate; H: High).

Table (3) represented studied main domain's items assessed, such that (Mean of score, Standard deviation, and Relative sufficiency) of likert score (Always, Sometimes, and Never) with scales assess (3, 2, and 1) respectively.

Most of studied items concerning of family environment main domain are assigned at a moderate level mostly, since they are accounted 9(81.81%), while only 2(18.18%) items were assessed at the low level, then followed mostly at a moderate level with respect to external environment main domain's items, since accounted 10(83.33%), while only 2(16.67%) items



were assessed at the low level, then followed mostly at a moderate level with respect to health problems related to vision, since accounted 9(69.23%), while only 4(16.67%) items were assessed at the low level, then followed mostly at a low level with respect to health problems related to musculoskeletal pain, since they are accounted 6(66.67%), while only 3(33.33%) items were assessed at the moderate level, then followed mostly at a moderate level by the psychological problems, since they are accounted 11(78.57%), while only 2(14.29%) items were assessed at the a high level, and only 1(4.17%) of them assessed at a low level.

Table 4: Simple Pearson's Correlation Coefficients among Different Responses of the Studied Domains

Simple Pearson's Correlation Coefficients Among studied Domains		External Environment	HPR to vision	HPR to Musculoskele tal pain	Psychological Problems
rho	Family Environment	0.503	0.152	0.165	0.414
	External Environment		0.176	0.234	0.423
	HPR to vision			0.351	0.250
	HPR to Musculoskeletal pain				0.259
P-value	Family Environment	0.000	0.000	0.000	0.000
	External Environment		0.000	0.000	0.000
	HPR to vision			0.000	0.000
	HPR to Musculoskeletal pain				0.000

HS: Highly Sig. At P<0.01; Statistical Hypothesis Is Based on Pearson’s Coefficient Test.

Table (4) shows in term simple correlation coefficients (Person's coefficients) of studied domains. There is a conventional significant level for extracted correlation coefficients among the studied domains in light of observing strong relationships at P<0.01 among all studied domains, and that indicated a meaningful limited interaction (The Covariance) that should be presented indeed.

Component's Matrix	Components	
	1	2
Family Environment Domain	0.834	
External Environment Domain	0.811	
Health problems related to vision Domain		0.822
Health problems related to Musculoskeletal pain Domain		0.790
Psychological Domain	0.696	
Initial Eigen values	2.196	1071
% of covariance	67.22	32.78
Suggested Named of extracted Factors	Psycho-Socio Impact Factor	Physical Impact Factor

Table5: Extracted Factors Matrix in Rotated Method with the Suggested Named for the Studied Grousted named "Psycho-Socio impact Factor, and Physical Impact Factor". That



extracted factors ordered in more powerful significant, with advantage at the first factor in (43.921%) of first total variance explained, then followed by second factor in (21.422%) of remaining total variance explained, and that led to be recorded (67.22%) of covariance explained for the first factor and (32.78%) of covariance explained for the second factor.

Discussion

In our study, the analysis Pearson's correlation coefficients aim to explore the interrelationships among different factors associated with Smartphone misuse. All the correlations presented in the table have a $P < 0.000$, indicating that these correlations are statistically highly significant ($P < 0.01$). The findings indicate a strong interconnectedness among various domains related to Smartphone misuse, this interconnectedness highlights the multifaceted impacts of Smartphone misuse on different aspects of an individual's life, from the family environment to health issues. pervious study reported, Students experienced sleep difficulties and visual changes, including dry eye disease, burning sensations in the eyes, conjunctival infection, reduced vision, and macular degeneration (Liu et al., 2022).. The studies on musculoskeletal disorders (MSDs) and Smartphone usage found that the most commonly affected body regions were the neck, followed by the shoulders, hands/wrists, and elbows in the general population (Mokhtarinia et al., 2024; Alotaibi et al., 2022; Soliman Elserty et al., 2020). Researchers discovered that Smartphone addicts experience detrimental effects on their interpersonal connections and mental health. This phenomenon can be ascribed to the fact that those who are too dependent on Smartphone s may have hindered their in-person social connections, resulting in a fear of missing out (FOMO) and anxiety. Ultimately, this can lead to feelings of loneliness and melancholy (Gökçearsan et al., 2021; Tangmunkongvorakul et al., 2020). Our study the results show of a factor analysis performed on the studied domains, which were extracted into two meaningful and significant interactions. The factors are named "Psycho-Socio Factor" and "Physical Impact Factor." The significance of the Psycho-Socio Factor indicates that the excessive use of Smartphone s has a substantial impact on family interactions and dynamics.

5. CONCLUSIONS

The observed correlations also indicate that there is a significant interconnection between studied main domains. This interconnection means that these factors do not operate in isolation from each other, but rather influence each other in the context of Smartphone use and impact. The analysis extracted two significant factors; the result of a factor analysis performed on data, first, has a suggested named "Psycho-Socio Impact Factor". This factor is the most significant, and explaining about 43.921% of the total variance. It includes variables related to family environment, external environment, and psychological domains. The second, factor has a suggested named "Physical Impacts Factor". This factor explains about 21.422% of the total variance and is associated with health problems related to vision and musculoskeletal pain, and that led to be recorded (67.22%) of covariance explained for the first factor and (32.78%) of covariance explained for the second factor.

Recommendation



1. Developing an instructional program with the goal of enhancing the quality of life for individuals suffering from Smartphone addiction, as well as holding workshops, courses, and conferences to increase awareness and information about the detrimental impacts of Smartphone addiction.
2. The researchers suggested that for future studies should focus on the phenomenon of acquired autism, particularly in children who are addicted to misuse Smartphone.

6. REFERENCES

1. Abdulkhaleq A Ali Ghalib, Baqer A, Munther S; (2012); "Health problems associated with the wrong use of computers in the city of Baghdad", Iraqi national journal of nursing specialties, vol.25, No.2.Achangwa, C., Ryu, H. S., Lee, J. K., & Jang, J.-D. (2022). Adverse Effects of Smartphone Addiction among University Students in South Korea: A Systematic Review. *Healthcare*, 11(1), 14. <https://doi.org/10.3390/healthcare11010014>
2. Alotaibi, M. S., Fox, M., Coman, R., Ratan, Z. A., & Hosseinzadeh, H. (2022). Smartphone Addiction Prevalence and Its Association on Academic Performance, Physical Health, and Mental Well-Being among University Students in Umm Al-Qura University (UQU), Saudi Arabia. *International Journal of Environmental Research and Public Health*, 19(6), 3710. <https://doi.org/10.3390/ijerph19063710>
3. Basheer, S. N., Arishi, T. M., Albar, N., Peeran, S. W., Peeran, S. A., & Zameer, M. (2021). Smartphone Addiction and Its Impact on Knowledge, Cognitive, and Psychomotor Skills of the Dental Students at Jazan University. *World Journal of Dentistry*, 11(6), 468–473. <https://doi.org/10.5005/jp-journals-10015-1778>
4. Cha, S.-S., & Seo, B.-K. (2018). Smartphone use and Smartphone addiction in middle school students in Korea: Prevalence, social networking service, and game use. *Health Psychology Open*, 5(1), 2055102918755046. <https://doi.org/10.1177/2055102918755046>
5. Gazzaz, Z. J., Baig, M., Al Alhendi, B. S. M., Al Suliman, M. M. O., Al Alhendi, A. S., Al-Grad, M. S. H., et al. (2018). Perceived stress, reasons for and sources of stress among medical students at Rabigh Medical College, King Abdulaziz University, Jeddah, Saudi Arabia. *BMC Med. Educ.* 18:29. <https://doi.org/10.1186/s12909-018-1133-2>
6. Gökçearsan, Ş., Yildiz Durak, H., Berikan, B., & Saritepeci, M. (2021). Smartphone Addiction, Loneliness, Narcissistic Personality, and Family Belonging Among University Students: A Path Analysis. *Social Science Quarterly*, 102(4), 1743–1760. <https://doi.org/10.1111/ssqu.12949>
7. Hassim, S. R., Arifin, W. N., Kueh, Y. C., & Yaacob, N. A. (2020). Confirmatory Factor Analysis of the Malay Version of the Smartphone Addiction Scale among Medical Students in Malaysia. *International Journal of Environmental Research and Public Health*, 17(11), 3820. <https://doi.org/10.3390/ijerph17113820>
8. Hundekari, J., Sisodiya, R., & Kot, L. (2021). Smartphone Vision Syndrome Associated with Prolonged Use of Digital Screen for Attending Online Classes during COVID-19 Pandemic among Medical Students: A Cross-sectional Study. *JOURNAL OF CLINICAL AND DIAGNOSTIC RESEARCH*. <https://doi.org/10.7860/jcdr/2021/46651.14399>



9. I.Mohammed Ali, S., & Nihad, M. (2021). Internet of Things for Education Field. *Journal of Physics: Conference Series*, 1897, 012076. <https://doi.org/10.1088/1742-6596/1897/1/012076>
10. Ismael, D. S. (2022). Smartphone overuse and vision problems among university students. *Zanco Journal of Pure and Applied Sciences*, 34(3). <https://www.iasj.net/iasj/article/237217>
11. Kee, I.-K., Byun, J.-S., Jung, J.-K., & Choi, J.-K. (2016). The presence of altered craniocervical posture and mobility in Smartphone -addicted teenagers with temporomandibular disorders. *Journal of Physical Therapy Science*, 28(2), 339–346. <https://doi.org/10.1589/jpts.28.339>
12. Kuorinka, I., Jonsson, B., Kilbom, A., Vinterberg, H., Biering-Sørensen, F., Andersson, G., & Jørgensen, K. (1987). Standardised Nordic questionnaires for the analysis of musculoskeletal symptoms. *Applied Ergonomics*, 18(3), 233–237. [https://doi.org/10.1016/0003-6870\(87\)90010-x](https://doi.org/10.1016/0003-6870(87)90010-x)
13. Kim, H., & Kim, H. (2013). Exercise rehabilitation for Smartphone addiction. *Journal of Exercise Rehabilitation*, 9(6), 500–505. <https://doi.org/10.12965/jer.130080>
14. Lee, M., Chung, S. J., Lee, Y., Park, S., Kwon, J.-G., Kim, D. J., Lee, D., & Choi, J.-S. (2020). Investigation of Correlated Internet and Smartphone Addiction in Adolescents: Copula Regression Analysis. *International Journal of Environmental Research and Public Health*, 17(16), 5806. <https://doi.org/10.3390/ijerph17165806>
15. Lee, J., & Cho, B. (2015). Effects of Self-Control and School Adjustment on Smartphone Addiction among Elementary School Students. *International Journal of Contents*, 11, 1–6. <https://doi.org/10.5392/IJoC.2015.11.3.001>
16. Liu, H., Zhou, Z., Huang, L., Zhu, E., Yu, L., & Zhang, M. (2022). Prevalence of Smartphone addiction and its effects on subhealth and insomnia: A cross-sectional study among medical students. *BMC Psychiatry*, 22(1), 305. <https://doi.org/10.1186/s12888-022-03956-6>
17. Liu, H., Zhou, Z., Zhu, E., Huang, L., & Zhang, M. (2022). Smartphone addiction and its associated factors among freshmen medical students in China: A cross-sectional study. *BMC Psychiatry*, 22(1), 308. <https://doi.org/10.1186/s12888-022-03957-5>
18. Mehammed-Ameen, O. H. R., Sameen, F. Y., & Hasan, N. N. (2018). Smartphone Addiction among Nursing College Students in Kirkuk University. *Kufa Journal for Nursing Sciences*, 8(2), Article 2. <https://doi.org/10.36321/kjns.vi20182.2261>
19. Mokhtarinia, H. R., Torkamani, M. H., Farmani, N., & Gabel, C. P. (2024). Smartphone addiction prevalence, patterns of use, and experienced musculoskeletal discomfort during the COVID-19 pandemic in a general Iranian population. *BMC Public Health*, 24(1), 161. <https://doi.org/10.1186/s12889-024-17654-3>
20. Nife, Nadia & Nihad, Marwah & Hassan, Alaa. (2022). The Internet of Things Privacy. https://www.researchgate.net/publication/364011915_The_Internet_of_Things_Privacy
21. Sarhan, A. (2024). Prevalence and Associated Factors of Smartphone Addiction Among Medical and Health Sciences University Students: A Cross-Sectional Study. *Palestinian Medical and Pharmaceutical Journal*, 9(1). <https://doi.org/10.59049/2790-0231.1097>
22. Shahrestanaki, E., Maajani, K., Safarpour, M., Ghahremanlou, H. H., Tiyuri, A., & Sahebkar, M. (2020). The Relationship between Smartphone Addiction and Quality of



- Life among Students at Tehran University of Medical Sciences. *Addicta: The Turkish Journal on Addictions*, 7(1). <https://doi.org/10.15805/addicta.2020.7.1.0080>
23. Sohn, S. Y., Rees, P., Wildridge, B., Kalk, N. J., & Carter, B. (2019). Prevalence of problematic Smartphone usage and associated mental health outcomes amongst children and young people: a systematic review, meta-analysis and GRADE of the evidence. *BMC psychiatry*, 19(1), 356. <https://doi.org/10.1186/s12888-019-2350-x>
 24. Soliman Elserty, N., Ahmed Helmy, N., & Mohamed Mounir, K. (2020). Smartphone addiction and its relation to musculoskeletal pain in Egyptian physical therapy students. *European Journal of Physiotherapy*, 22(2). <https://doi.org/10.1080/21679169.2018.1546337>
 25. Tangmunkongvorakul, A., Musumari, P. M., Tsubohara, Y., Ayood, P., Srithanaviboonchai, K., Techasrivichien, T., Suguimoto, S. P., Ono-Kihara, M., & Kihara, M. (2020). Factors associated with Smartphone addiction: A comparative study between Japanese and Thai high school students. *PLOS ONE*, 15(9), e0238459. <https://doi.org/10.1371/journal.pone.0238459>
 26. Twenge, J. M., Joiner, T. E., Rogers, M. L., & Martin, G. N. (2018). Increases in Depressive Symptoms, Suicide-Related Outcomes, and Suicide Rates Among U.S. Adolescents After 2010 and Links to Increased New Media Screen Time. *Clinical Psychological Science*, 6(1), 3–17. <https://doi.org/10.1177/2167702617723376>
 27. Wang, J., Hao, Q.-H., Peng, W., Tu, Y., Zhang, L., & Zhu, T.-M. (2023). Relationship between Smartphone addiction and eating disorders and lifestyle among Chinese college students. *Frontiers in Public Health*, 11, 1111477. <https://doi.org/10.3389/fpubh.2023.1111477>