
Effect of Gender Difference on Left Ventricular and Left Atrial Parameters in Patients with Essential Hypertension

Dhuha Qasim Mohammed^{1*}, Asaad Hasan Noaman Al-Aboodi²

^{1*}Department of Physiology, Faculty of Medicine, University of Kufa, Iraq.

²Department of Physiology, Faculty of Medicine, University of Kufa. Al-Furat Teaching Hospital, Najaf, Iraq.

Email: ²asaadh.alaboodi@uokufa.edu.iq

Corresponding Email: ^{1*}dhuha.qasim14@gmail.com

Received: 29 March 2024

Accepted: 17 June 2024

Published: 01 August 2024

Abstract: *Background: arterial hypertension has a high impact on the heart including structural and functional changes, these changes may differ between males and females according to multiple causes. Aim: Establish if there is any difference in LV and LA parameters between males and females with hypertension. Patient and Methods: cross-sectional analytic study. The study samples were obtained from the Al-Furat Teaching Hospital's Echocardiographic Consultation Unit in the Al-Najaf Governorate between 1st September 2023, and 20th April 2024. Sixty-eight known cases of hypertensive patients with age of (18-60 years old). All patients are males and females. Clinical evaluation, anthropometric assessments, Doppler echocardiography were done for each one of them to assess the left atrial volumes and function and LV parameters. Results: the hypertensive patients show a significant difference in accordance with weight, height, age, and BSA (all $P<0.05$). at the same time IVS, LVPW and LVM shows a significant difference between males and females (all $P<0.05$). LA parameters (LA passive volume and conduit function) also shows a significant difference between two groups ($P<0.05$). conclusion: LV parameters affected by gender difference while LA parameters affected by age, ethnicity, type of treatment and compliance to it rather than the gender difference.*

Keywords: Gender Difference, LV Parameters, LA Parameters, Hypertension.

1. INTRODUCTION.

Arterial hypertension causes structural and functional changes in the heart, collectively named hypertensive heart disease (1). The heart is normally smaller in females than in males from puberty onwards (2), largely due to differences in body size and composition (3). Current



guidelines therefore recommend sex-specific threshold values for optimal detection of hypertensive heart disease by echocardiography (4). The hallmark of hypertensive heart disease and a potent prognostic indicator in hypertension is left ventricular hypertrophy, or LVH. In females compared to males, hypertension LVH is more common and less treatable with antihypertensive medication (5). Regardless of attained blood pressure levels, persistent LVH is specifically linked to elevated arterial stiffness, an increased risk of CV events, and mortality during follow-up. An additional prevalent indicator of hypertension heart disease is a dilated left atrium (LA). Increased CVD, specifically atrial fibrillation, heart failure (HF), and ischemic stroke, is linked to LA dilatation (6, 7). Another study reported that in middle-aged subjects with obesity without known CVD, LA dilatation was significantly more prevalent in females than in males, and particularly associated with co-presence of hypertension and

increased arterial stiffness (9). Urbina, Ferranti (15) stated that the hemodynamic load has a major influence on left ventricular features, including cardiac remodeling, because males and females have different blood pressure variations and hemodynamic loads. For instance, males with hypertension usually have greater systolic blood pressure than females do, which may lead to greater ventricular hypertrophy and higher LVM. According to Dobrowolski, Januszewicz (14) state that variations in how men and women respond to antihypertensive medications and other forms of treatment may have an impact on the structure and functionality of the heart.

2. RELATED WORKS

The results of Gerdt and colleagues showed that females exhibit higher LV myocardial function and ejection fraction than males, independent of LV geometry (5). Another study by Iosi and colleagues showed that LA dilatation is more common in females. In older subjects with hypertension and LVH, LA dilatation was significantly more prevalent in females (8). Another study reported that in middle-aged subjects with obesity without known CVD, LA dilatation was significantly more prevalent in females than in males, and particularly associated with co-presence of hypertension and increased arterial stiffness (9). Urbina, Ferranti (15) stated that the hemodynamic load has a major influence on left ventricular features, including cardiac remodeling, because males and females have different blood pressure variations and hemodynamic loads. For instance, males with hypertension usually have greater systolic blood pressure than females do, which may lead to greater ventricular hypertrophy and higher LVM. According to Dobrowolski, Januszewicz (14) state that variations in how men and women respond to antihypertensive medications and other forms of treatment may have an impact on the structure and functionality of the heart. Compliance with therapies and efficacy differences between the sexes may be the cause of differences in LV mass and dimensions. Kunišek and Kunišek (17) found that there were larger heart cavities and LVM in the male participants. These gender differences vanished when body surface area (LVMI) was used to index LVM. In terms of obesity, 38% of women and 26% of men were affected. The outcomes verified that obesity has an impact on LVM. The LA parameters are not significantly affected by gender ($p > 0.05$) due to the following reasons. Badano, Miglioranza (18) who studied



and increasing arterial stiffness, both of which have a role in the development of hypertension (11). Those who are taller typically have higher blood pressure than those who are shorter. This association is explained by the fact that in taller individuals, there is a greater hydrostatic pressure generated between the heart and the brain, necessitating higher blood pressure in order to provide appropriate cerebral perfusion. A higher body mass index

3. METHODOLOGY

3.1. Study Population:

This study uses cross-sectional analysis. The study samples were collected between September 1, 2023, and April 20, 2024, from the Echocardiographic Consultation Unit of Al-Furat Teaching Hospital in the Al-Najaf Governorate. There are documented cases of hypertension patients between the ages of 18 and 60. There are both male and female patients. This study excluded patients with diabetes mellitus, coronary artery disease, significant valvular disease, a history of atrial or ventricular arrhythmias, prior cardiac surgery, an implanted device, or any other chronic diseases (e.g., chronic kidney failure, chronic liver disease, etc.). The same physician collected the echocardiographic parameters apart from the clinical information. With SPSS V21, the statistical analysis is performed. All subjects had their ECGs and anthropometric measurements (weight and height) taken. Each patient's body surface area (BSA) was computed.

3.2. Echocardiography:

Every individual had a pulsed wave Doppler and 2D vivid E9 echocardiogram conducted utilizing a transducer that was connected to the ultrasound equipment. The echocardiography evaluation was conducted while a continuous three-lead ECG was being obtained. The American Society of Echocardiography's recommendations have been adhered to for conventional Doppler echocardiographic examinations (8). To acquire a four-chamber apical image, the transducer was positioned with the indicator pointing toward the left flank at the apex beat level at the fifth intercostal space at the mid-clavicular line. As a result, measurements were made of LAMax, LAMin, pre-atrial volume, E/A, DT, and E/é. The following parameters were measured using a parasternal long-axis view: IVS, LVEDV, and LVPW. The transducer was positioned between the nipples with the indication pointing toward the right shoulder. Following the guidelines provided by the American Society of Echography (ASE), the left ventricular mass (LVM) was measured and computed using Devereux's formula. A body surface area indexed LVM (LVMI) of greater than 115 g/m² in men and greater than 95 g/m² in women was considered left ventricular hypertrophy (LVH). It was feasible to differentiate between the concentric LVH if RWT \geq 0.42 and the eccentric LVH if RWT < 0.42 based on the relative wall thickness (RWT), which was determined by the 2 LVPW/LVEDV.

3.2. Statistical Analysis:

In this cross-sectional study, statistical analysis was performed using the statistical package SPSS for Windows (version 21, SPSS Inc., Chicago, IL, USA). Data analyzed as per group I for females (total: n=40) compared with group II for males (total: n=28) were shown as mean



± SD. Continuous variables were compared using independent sample t-tests. A P value < 0.05 was adopted to indicate statistical significance (10).

4. RESULTS AND 4. DISCUSSION

4.1. Demographic Characteristics:

Demographic characteristics of the population are given in Table 1. The mean age of subjects was 47.85 years and the results show a significant difference in age between the hypertensive people (p=0.00001). weight, height, and BSA also show a significant difference between hypertensive people (p= 0.002, 0.0003, and 0.000001 respectively).

Table 1: Demographic data

SD: standard deviation, **BSA:** body surface area, **Symbol *:** significant.

Demographic data	Hypertensive patients (n=68) Mean ± SD	P- value
Age (years)	47.85±8.95	0.00001*
Height (cm)	164.26±8.74	0.0003*
Weight (kg)	87.94±15.68	0.0020*
BSA	1.99±0.20	0.000001*

On the other hand, Table 2 revealed that mean of IVS in male group (11.39±1.449) was higher than the mean of female group (10.15±1.511), so there was a significant increase (p<0.05) in IVS thickness in the male group compared with the female group. The mean of LVPW thickness in the male group (10.93±1.245) was higher than the female group (10.03±1.561), so there was a significant increase (p<0.05) in LVPW thickness in the male group compared with the female group. The mean of relative wall thickness (RWT) of left ventricle in the male and the female group were (0.50±0.06) and (0.46±0.09), which was not significantly different between the two groups (p>0.05). The mean of LVMI in the male and female group were (104.31±23.49) and (96.44 ±20.89), which was not significantly different between the two groups (p>0.05). The mean of LVM was (219.43 ±54.25) in male group higher significantly than the mean of LVM (185.69± 45.83) in female group, (p<0.05).

Table (2): LV parameters according to gender specification.

LV: left ventricle, **LVM:** left ventricular mass, **LVMI:** left ventricular mass index, **SD:** standard deviation, **IVS:** interventricular septum, **LVPW:** left ventricular posterior wall, **RWT:** Relative wall thickness, **Symbol *:** significant.

LV parameters	Female group (n=40) mean±SD	Male group (n=28) mean±SD	P- value
IVS	10.15±1.511	11.39±1.449	0.001*
LVPW	10.03±1.561	10.93±1.245	0.013*
RWT	0.46 ±0.09	0.50± 0.06	0.109
LVMi	96.44 ±20.89	104.31±23.49	0.151
LVM	185.69 ± 45.83	219.43 ±54.25	0.007*



Table 3 shows all volumes were not significantly different between male and female groups ($p > 0.05$). Conduit function was higher significantly in males than in females ($p < 0.05$).

Table (3): LA parameters related to gender specification.
LA: left atrium, **SD:** standard deviation, **symbol *:** significant.

LA parameters	Female group (n=40) mean±SD	Male group (n=28) mean±SD	P- value
Maximum volume	57.43±10.107	62.21±16.231	0.139
Minimum volume	16.00±5.729	15.79±5.750	0.880
Pre P volume	30.80±8.392	29.00±7.921	0.376
Reservoir volume	41.43±8.221	46.43±13.287	0.060
LA contractile volume	14.80±5.766	13.21±6.286	0.286
Passive emptying volume	26.63±8.095	33.21±11.571	0.007*
Reservoir function	41.43±8.221	46.43±13.287	0.060
Conduit function	26.63±8.095	33.21±11.571	0.007*
Booster function	14.80±5.766	13.21±6.286	0.286
Passive emptying function	56.88 ±10.11	61.74 ±16.24	0.134
Active emptying function	30.27 ±8.400	28.44 ±7.96	0.369
Total emptying fraction (EF)	57.14 ±10.09	61.95 ±16.23	0.137

Discussion

The demographic data of our study differ significantly in hypertensive patients and this means that these parameters play an important role in developing hypertension. Age-related physiological changes include endothelial dysfunction and increasing arterial stiffness, both of which have a role in the development of hypertension (11). Those who are taller typically have higher blood pressure than those who are shorter. This association is explained by the fact that in taller individuals, there is a greater hydrostatic pressure generated between the heart and the brain, necessitating higher blood pressure in order to provide appropriate cerebral perfusion. A higher body mass index (BMI) and body weight are linked to a higher risk of hypertension. Being overweight causes a number of physiological alterations, including insulin resistance, endothelial dysfunction, and an increase in sympathetic nervous system activity, all of which have a role in the development of hypertension (12). BSA has a positive correlation with blood pressure and is determined using height and weight. Blood pressure is often greater in people with larger BSAs than in people with smaller BSAs (13).

Table (2) shows that the association demonstrated a significant difference ($p < 0.05$) in IVS, LVPW, and LVM between male and female hypertensive patients. Compared to females, males had higher values of these factors. This clarifies how the values for these factors differ in men and women on average. As a result, women's LV reacts differently to continuous pressure overloaded in hypertension. (14-16) explain the reason of this. According to Vitale,



Mendelsohn (16) The form and function of the cardiovascular system can be influenced by the levels of estrogen and testosterone, two hormones that are different in men and women. For example, vasodilation and the inhibition of cardiac hypertrophy have been associated with the cardioprotective effects of estrogen, which could account for the differences in LVPW, LVM, and IVS between the sexes. Urbina, Ferranti (15) stated that the hemodynamic load has a major influence on left ventricular features, including cardiac remodeling, because males and females have different blood pressure variations and hemodynamic loads. For instance, males with hypertension usually have greater systolic blood pressure than females do, which may lead to greater ventricular hypertrophy and higher LVM. According to Dobrowolski, Januszewicz (14) state that variations in how men and women respond to antihypertensive medications and other forms of treatment may have an impact on the structure and functionality of the heart. Compliance with therapies and efficacy differences between the sexes may be the cause of differences in LV mass and dimensions. Kunišek and Kunišek (17) found that there were larger heart cavities and LVM in the male participants. These gender differences vanished when body surface area (LVMI) was used to index LVM. In terms of obesity, 38% of women and 26% of men were affected. The outcomes verified that obesity has an impact on LVM. The LA parameters are not significantly affected by gender ($p>0.05$) due to the following reasons. Badano, Miglioranza (18) who studied the effect of gender difference on LA parameters and suggested that although women have lower diastolic compliance and higher systolic chamber function than men do, there are gender differences in left ventricular chamber function as well, although these differences do not directly affect the LA parameters. Research has demonstrated that left atrial volumes, irrespective of gender, rise markedly with age. Doria de Vasconcellos, Win (19) explained that although age and ethnicity have been taken into consideration when establishing reference values for left atrial volumes, emptying fractions, these data do not show a substantial influence of gender on LA parameters.

5. CONCLUSION

there is a significant difference in LV parameters between males and females while this difference doesn't find in LA parameters except of conduit function and passive emptying volume, the reason behind this, is these parameters affected by age, ethnicity, compliance to the treatment of hypertension and the type of treatment rather than the gender while conduit function and passive emptying volume depend directly on the left ventricular filling pressure. estrogen, which could account for the differences in LVPW, LVM, and IVS between the sexes. Urbina, Ferranti (15) stated that the hemodynamic load has a major influence on left ventricular features, including cardiac remodeling, because males and females have different blood pressure variations and hemodynamic loads. For instance, males with hypertension usually have greater systolic blood pressure than females do, which may lead to greater ventricular hypertrophy and higher LVM. According to Dobrowolski, Januszewicz (14) state that variations in how men and women respond to antihypertensive medications and other forms of treatment may have an impact on the structure and functionality of the heart. Compliance with therapies and efficacy differences between the sexes may be the cause of differences in LV mass and dimensions. Kunišek and Kunišek (17) found that there were larger heart cavities and



LVM in the male participants. These gender differences vanished when body surface area (LVMI) was used to index LVM. In terms of obesity, 38% of women and 26%

Ethics Approval and Consent to Participate

The Department of Physiology, Faculty of Medicine, University of Kufa and department of Echocardiography, Al-Furat Teaching Hospital, Najaf, Iraq, authorized the study design, and all patients had provided informed consent.

Acknowledgment

I would want to express my gratitude to the Al-Furat Teaching Hospital's echocardiography unit team for their assistance and cooperation in getting the study samples.

6. REFERENCES

1. Whelton PK, Carey RM, Aronow WS, Casey DE, Jr., Collins KJ, Dennison Himmelfarb C, et al. 2017 ACC/AHA/AAPA/ABC/ACPM/AGS/APhA/ASH/ASPC/NMA/PCNA Guideline for the Prevention, Detection, Evaluation, and Management of High Blood Pressure in Adults: Executive Summary: A Report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines. *Hypertension*. 2018;71(6):1269-324.
2. de Simone G, Devereux RB, Daniels SR, Meyer RA. Gender differences in left ventricular growth. *Hypertension*. 1995;26(6 Pt 1):979-83.
3. De Simone G, Devereux RB, Chinali M, Roman MJ, Barac A, Panza JA, et al. Sex differences in obesity-related changes in left ventricular morphology: the Strong Heart Study. *J Hypertens*. 2011;29(7):1431-8.
4. Yang BY, Qian Z, Howard SW, Vaughn MG, Fan SJ, Liu KK, et al. Global association between ambient air pollution and blood pressure: A systematic review and meta-analysis. *Environ Pollut*. 2018;235:576-88.
5. Gerds E, Okin PM, Simone Gd, Cramariuc D, Wachtell K, Boman K, et al. Gender Differences in Left Ventricular Structure and Function During Antihypertensive Treatment. *Hypertension*. 2008;51(4):1109-14.
6. Gupta S, Matulevicius SA, Ayers CR, Berry JD, Patel PC, Markham DW, et al. Left atrial structure and function and clinical outcomes in the general population. *Eur Heart J*. 2013;34(4):278-85.
7. Sardana M, Lessard D, Tsao CW, Parikh NI, Barton BA, Nah G, et al. Association of Left Atrial Function Index with Atrial Fibrillation and Cardiovascular Disease: The Framingham Offspring Study. *J Am Heart Assoc*. 2018;7(7).
8. Losi MA, Mancusi C, Midtbø H, Saeed S, de Simone G, Gerds E. Impact of estimated left atrial volume on prognosis in patients with asymptomatic mild to moderate aortic valve stenosis. *Int J Cardiol*. 2019;297:121-5.
9. Halland H, Lønnebakken MT, Pristaj N, Saeed S, Midtbø H, Einarsen E, et al. Sex differences in subclinical cardiac disease in overweight and obesity (the FATCOR study). *Nutr Metab Cardiovasc Dis*. 2018;28(10):1054-60.



10. WAYNE W. DANIEL PDCLC, PH.D., PSTAT. Biostatistics A Foundation for Analysis in the Health Sciences. 2018.
11. Evans JM, Wang S, Greb C, Kostas V, Knapp CF, Zhang Q, et al. Body Size Predicts Cardiac and Vascular Resistance Effects on Men's and Women's Blood Pressure. *Front Physiol.* 2017;8:561.
12. Devonshire AL, Hager ER, Black MM, Diener-West M, Tilton N, Snitker S. Elevated blood pressure in adolescent girls: correlation to body size and composition. *BMC Public Health.* 2016;16(1):78.
13. Raphadu TT, Staden MV, Dibakwane WM, Monyekei KD. A Non-Invasive Investigation into the Prevalence of Higher than Normal Blood Pressure, Hypertension and the Association between Blood Pressure and Body Weight in Male and Female Adolescents in the Polokwane Local Municipality, Limpopo-South Africa: A Cross-Sectional Study. *Children (Basel).* 2020;7(3).
14. Dobrowolski P, Januszewicz M, Klisiewicz A, Prejbisz A, Warchol-Celinska E, Michalowska I, et al. Echocardiographic assessment of left ventricular morphology and function in patients with fibromuscular dysplasia: the ARCADIA-POL study. *Journal of Hypertension.* 2018;36(6):1318-25.
15. Urbina EM, Ferranti Sd, Steinberger J. Observational Studies May Be More Important Than Randomized Clinical Trials. *Hypertension.* 2014;63(4):638-40.
16. Vitale C, Mendelsohn ME, Rosano GMC. Gender differences in the cardiovascular effect of sex hormones. *Nature Reviews Cardiology.* 2009;6(8):532-42.
17. Kunišek J, Kunišek L. IMPACT OF BLOOD PRESSURE COMPONENTS ON LEFT VENTRICULAR HYPERTROPHY REMODELING. *Acta Clin Croat.* 2018;57(4):638-45.
18. Badano LP, Miglioranza MH, Mihaila S, Peluso D, Xhaxho J, Marra MP, et al. Left Atrial Volumes and Function by Three-Dimensional Echocardiography: Reference Values, Accuracy, Reproducibility, and Comparison With Two-Dimensional Echocardiographic Measurements. *Circ Cardiovasc Imaging.* 2016;9(7).
19. Doria de Vasconcellos H, Win TT, Chamera E, Hong SY, Venkatesh BA, Young P, et al. References Values for Left Atrial Volumes, Emptying Fractions, Strains, and Strain Rates and Their Determinants by Age, Gender, and Ethnicity: The Multiethnic Study of Atherosclerosis (MESA). *Acad Radiol.* 2021;28(3):356-63.