
Effect of Ageing Process on Diastolic Function of the Heart

Fatima F. Al-Muhanna^{1*}, Amina A. B. Al-Dejeli²

^{1*}Master student, College of medicine, University of Kufa, Najaf, Iraq.

²Department of physiology, Faculty of medicine, University of Kufa. Department of Echo Laboratory, Al-Najaf Center for Cardiac Surgery and Transcatheter Therapy/ Al-Sader Teaching Hospital, Iraq.

Corresponding Email: ^{1*}FatimaF.A114@gmail.com

Received: 29 March 2024

Accepted: 17 June 2024

Published: 01 August 2024

Abstract: *As people age, diastolic function deteriorates. (1). Aging is a significant risk factor for the emergence of structural and functional problems in the heart (2). Diastolic dysfunction is a significant problem observed with age. Echocardiography is used to investigate the presence of diastolic dysfunction, and the pathophysiologic factors affecting left ventricular and left atrial anatomy and function. (3)*

Aim of the study: *The objective of this study is to investigate the impact of ageing process on diastolic function of the heart which is assessed by echocardiography.*

Methods: *in our cross sectional study, cases were collected from Al-Najaf Center for Cardiac Surgery and Transcatheter Therapy/ Al-Sader Teaching Hospital. It included 50 patients aged 38-86 who underwent 2d echocardiography to assess diastolic function of the heart.*

Results: *The results of the study obviously showed that left ventricular diastolic function steadily declines with age (P value < 0.05).*

Conclusion: *left ventricular diastolic function of the heart significantly declines with age.*

Keywords: *Diastolic Dysfunction, Age, Echocardiography, Cardiac Remodelling.*

1. INTRODUCTION

Diastolic dysfunction (DD) is a general word for a number of physiological mechanisms that enable the left ventricle, or LV, to be filled with enough blood to meet the body's current needs while maintaining just enough pressure to avoid pulmonary congestion. (4)

Causes of diastolic dysfunction include the following: (5)

- Hypertrophy of the left ventricle
- Ischemia



- Pericardial disease
- Remodeling and fibrosis (post myocardial infarction or in chronic heart failure)
- Age > 65 years

Heart remodeling, particularly left ventricular (LV) remodeling, is defined as a complex alteration in the structure and function of the heart that advances with the age of the patient (6), and is linked to an increased risk of cardiovascular morbidity and mortality (7). Therefore, it is critical to clarify the mechanisms of LV remodeling in order to provide medical care for the aged people. Numerous studies have demonstrated a correlation between ageing and LV remodeling ; however, it is unclear whether this is a direct or indirect relationship because ageing is also associated with the development and aggravation of a number of diseases that impact structural/functional LV remodeling, including hypertension, obesity, diabetes, chronic kidney disease, and hyperlipidemia.(6)

Echocardiography is essential for assessment of left ventricular diastolic function. According to the most recent guidelines published by the European Society of Cardiology, echocardiography is a Class I Level C recommendation because of its safety, availability, and accuracy in the diagnosis and treatment of heart failure. (8).

According to current guidelines for Doppler echocardiography, diastolic function should be assessed via early to late diastolic transmitral flow velocity (E/A), and left ventricular filling pressure should be estimated using E to early diastolic mitral annular tissue velocity (E/e'). (4) E/A ratio < 0.8 has been identified as a straightforward yet powerful biomarker of LVDD. (9)

level of structural integrity, there is an increase in connective tissue matrix content (16-18), a decrease in myocyte number (17), and an increase in myocyte size (18). Apoptosis and cell necrosis produce a decrease in the quantity of myocytes (19).

2. RELATED WORKS

The results of Dugo and colleagues about assessment of diastolic function by echocardiography in elderly people, showed that diastolic function of the left ventricle significantly decrease with advanced age. (8) in another study by Meluzin et al, about diastolic dysfunction in elderly patients , they showed that The "physiological" diastolic dysfunction in the aged can be explained by a decline in passive left ventricular filling qualities and a worsening of left ventricular relaxation. (10) another study reported that an early slowing of LV relaxation and diastolic suction begins in early middle age, with the greatest drop reported in elderly people, and a decrease in ventricular relaxation is likely responsible for the changes in LV diastolic filling with senescence. (11)

Essential for assessment of left ventricular diastolic function. According to the most recent guidelines published by the European Society of Cardiology, echocardiography is a Class I Level C recommendation because of its safety, availability, and accuracy in the diagnosis and treatment of heart failure

Numerous studies have demonstrated a correlation between ageing and LV remodeling; however, it is unclear whether this is a



The most frequent pathologic cause of elevated afterload is arterial hypertension, which results in left ventricular (LV) hypertrophy (LVH) and wall thickening in order to maintain tissue perfusion and pump function. LVH is closely linked to preclinical cardiovascular abnormalities, elevated cardiovascular morbidity and mortality

3. METHODOLOGY

Study sample: The study population consists of patients with negative coronary artery disease (CAD) confirmed by percutaneous coronary intervention (PCI) who are participating in a population-based cross-sectional study. They were taken from the echocardiology department of Al-Najaf Center for Cardiac Surgery and Transcatheter Therapy/Al-Sader Teaching Hospital. The study included 50 patients ranging in age from 38 to 86 years. They all underwent 2D echocardiography to evaluate diastolic function. We only recruited patients with sinus rhythm, a left ventricular ejection fraction of 55% or more, no valvular disease, no evidence of pulmonary hypertension, negative CAD confirmed by PCI, and with no right ventricular dysfunction.

Every participant was informed about the study's objectives, procedures, and methods, considering the option to participate or not. They gave express and willing consent. The study was also approved by the ethical committee of University of Kufa, Faculty of Medicine.

Echocardiographic Assessment:

Each participant was submitted to 2D echocardiographic assessment for systolic and diastolic function. Regarding diastolic function, patients were categorized into Grade 0 which is normal, grade I which is impaired relaxation, grade II that is pseudonormal, grade III restrictive and grade IV diastolic dysfunction irreversibly restrictive. (12) In this study, early to late diastolic transmitral flow velocity (E/A) have been used for assessment of left ventricular diastolic function. The machine used for echocardiographic assessment was GE Vivid E9 Ultrasound System which was a high performance digital ultrasound imaging system.

Statistical Analysis:

The data was analyzed using version 26 of the Statistical Package for Social Science (SPSS) software. Categorical variables were expressed as frequencies and percentages, and the Chi-square test have been used to measure their level of significance and correlations. The P value of 0.05 was employed in this study. P values < or =0.05 were considered significant.

4. RESULTS AND DISCUSSION

Table1 lists the frequency of Demographic Characteristics and diastolic function of the patients. Table2 lists the relationship between age and the patients' diastolic function. Table 3. Shows the relationship between gender and the patients' diastolic function and Table 4. Shows the relationship between HTN and the patients' diastolic function.

The study's findings clearly indicate a steady decline in left ventricular diastolic function with aging, (P value = 0.01). it also found a positive correlation between HTN and DD (p value= 0.03). There is no significant relationship between gender and diastolic function (p value = 0.1)

Table 1. Demographic Characteristics and risk factors of the patients.

	30-49	5	10%	68.2 ± 1.6
	50-69	29	58%	59.55 ± 11.04
	70-89	16	32%	63.88 ± 10.8
	total	50	100%	61.8 ± 10.7
	Male	33	66%	
	female	17	34%	
	Hypertensive	41	82%	
	Non-hypertensive	9	18%	
	Grade 0	2	4%	
	Grade I	22	44%	
	Grade II	16	32%	
	Grade III	10	20%	

Table 2. The relationship between age and the patients' diastolic function

age	Diastolic function				total	P value
	Grade 0	Grade 1	Grade 2	Grade 3		
30-49	2 (4%)	3 (6%)	0 (0%)	0 (0%)	5 (10%)	0.001
50-69	0 (0%)	14 (28%)	10 (20%)	5 (10%)	29 (58%)	
70-89	0 (0%)	5 (10%)	6 (12%)	5 (10%)	16 (32%)	
total	2 (4%)	22 (44%)	16 (32%)	10 (20%)	50 (100%)	

Table 3. The relationship between gender and the patients' diastolic function

gender	Diastolic function				total	P value
	Grade 0	Grade 1	Grade 2	Grade 3		
male	2 (4%)	17 (34%)	7 (14%)	7 (14%)	33 (66%)	0.1
female	0 (0%)	5 (10%)	9 (18%)	3(6%)	17 (14%)	
total	2 (4%)	22 (44%)	16 (32%)	10 (20%)	50 (100%)	

Table 4. The relationship between HTN and the patients' diastolic function

HTN	Diastolic function				total	P value
	Grade 0	Grade 1	Grade 2	Grade 3		
hypertensive	2 (4%)	13 (26%)	16 (32%)	10 (20%)	41(82%)	0.03
Non-hypertensive	0 (0%)	9 (18%)	0 (0%)	0 (0%)	9 (18%)	
total	2 (4%)	22 (44%)	16 (32%)	10 (20%)	50 (100%)	

Regarding age, our finding was consistent with a number of earlier studies. (1, 8, 13-15) Toba and colleagues stated that Regardless of LV hypertrophy, age was independently connected



with LV concentric/functional alterations, indicating that aging is independently engaged in the development of LV remodelling. (6). Another study showed that Age considerably reduced the E/A ratio ($p < 0.001$), whereas gender had no effect ($p = 0.298$). E/e ratio was higher in women than in males ($p < 0.001$) and increased considerably with age ($p < 0.001$). (13).

On the level of structural integrity, there is an increase in connective tissue matrix content (16-18), a decrease in myocyte number (17), and an increase in myocyte size (18). Apoptosis and cell necrosis produce a decrease in the quantity of myocytes (19). Fibroblasts replace lost myocytes, causing the remaining myocytes to enlarge. Interstitial fibrosis results from the fibroblasts' production of collagen, which makes the heart less flexible and more rigid (20). Both systolic contraction and diastolic relaxation are impacted by the less flexible and stiffer ventricle. (21) Systolic contraction time prolongation and left ventricular hypertrophy (LVH) are caused by persistently high afterload (22) by rigid vasculature. In turn, early diastole is impacted by prolonged systolic contraction. (21)

Regarding hypertension, Bamaiyi concluded that Left ventricular mass index (LVMI) and relative wall thickness (RWT) only contribute a small amount to the effect of BP on diastolic function, despite the fact that structural LV remodelling is independently linked to alterations in LV diastolic performance. Therefore, it is likely that the majority of BP-associated reductions in LV diastolic performance are a transition phase unrelated to LV hypertrophy or concentric remodelling. (23) The most frequent pathologic cause of elevated afterload is arterial hypertension, which results in left ventricular (LV) hypertrophy (LVH) and wall thickening in order to maintain tissue perfusion and pump function. LVH is closely linked to preclinical cardiovascular abnormalities, elevated cardiovascular morbidity and mortality. (24) According to a study done by Tran et al., adolescents with increased blood pressure are at risk for subclinical alterations in their systolic and diastolic cardiac function. (25) According to Another study, around 48% of patients with newly diagnosed but untreated hypertension exhibit abnormalities in their left ventricle's diastolic function. (26)

Regarding gender, our study had shown no difference between male and female regarding diastolic function, $P > 0.05$. Shim and colleagues in their study about gender difference regarding the effect of arterial stiffness on the diastolic function of the heart, stated that the measurements of arterial stiffness and LV diastolic function strongly correlate only in females. (27) Carvalho and colleagues also had a result different from ours. Their study showed that even with normal aging, women's diastolic function declines more quickly than men's. (13) This difference between results may be caused by the huge difference in sample size as their study included 14,298 patients, and difference in the age group as the ages of the two oldest and youngest patients were 98 and 7, respectively.

5. CONCLUSION

This study found that left ventricular diastolic function of the heart significantly declines with age. Hypertension, Bamaiyi concluded that Left ventricular mass index (LVMI) and relative wall thickness (RWT) only contribute a small amount to the effect of BP on diastolic



function, despite the fact that structural LV remodelling is independently linked to alterations in LV diastolic performance. Therefore, it is likely that the majority of BP-associated reductions in LV diastolic performance are a transition phase unrelated to LV hypertrophy or concentric remodelling. (23) The most frequent pathologic cause of elevated afterload is arterial hypertension, which results in left ventricular (LV) hypertrophy (LVH) and wall thickening in order to maintain tissue perfusion and pump function. LVH is closely linked to preclinical cardiovascular abnormalities, elevated cardiovascular morbidity and mortality. (24) According to a study done by Tran et al., adolescents with increased blood pressure are at risk for subclinical alterations in their systolic and diastolic cardiac function. (25) According to Another study, around 48% of patients with newly diagnosed but untreated hypertension exhibit abnormalities in their left ventricle's diastolic function. (26)

6. REFERENCES

1. Santoro A, Alvino F, Antonelli G, Cassano FE, De Vito R, Cameli M, et al. Age related diastolic function in amateur athletes. *The International Journal of Cardiovascular Imaging*. 2015;31(3):567-73.
2. Saheera S, Krishnamurthy P. Cardiovascular Changes Associated with Hypertensive Heart Disease and Aging. *Cell Transplantation*. 2020;29:0963689720920830.
3. Nagueh Sherif F. Left Ventricular Diastolic Function. *JACC: Cardiovascular Imaging*. 2020;13(1_Part_2):228-44.
4. Mitter SS, Shah SJ, Thomas JD. A test in context: E/A and E/e' to assess diastolic dysfunction and LV filling pressure. *Journal of the American College of Cardiology*. 2017;69(11):1451-64.
5. Groban L, Garner CR. 3 - Diastolic Function. In: Groban L, Garner CR, editors. *TEE Pocket Manual (Second Edition)*: Elsevier; 2018. p. 23-38.
6. Toba A, Kariya T, Aoyama R, Ishiyama T, Tsuboko Y, Takeda K, et al. Impact of age on left ventricular geometry and diastolic function in elderly patients with treated hypertension. *Blood Pressure*. 2017;26(5):264-71.
7. Wu Q-Q, Xiao Y, Yuan Y, Ma Z-G, Liao H-H, Liu C, et al. Mechanisms contributing to cardiac remodelling. *Clinical Science*. 2017;131(18):2319-45.
8. Dugo C, Rigolli M, Rossi A, Whalley GA. Assessment and impact of diastolic function by echocardiography in elderly patients. *J Geriatr Cardiol*. 2016;13(3):252-60.
9. Johansson B, Lundin F, Tegerback R, Boj  L. E/a' ratio a simple detector of left ventricular dysfunction in patients with decreased ejection fraction. *Scand Cardiovasc J*. 2018;52(1):20-7.
10. Meluzin J, Podrouz kov H, Gregorov Z, Panovsk R. [Diastolic dysfunction in the elderly subjects. Disease or a physiological manifestation of ageing?]. *Vnitřni lékařství*. 2013;59 5:392-6.
11. Carrick-Ranson G, Carrick-Ranson G, Hastings JL, Bhella PS, Shibata S, Shibata S, et al. Effect of healthy aging on left ventricular relaxation and diastolic suction. *American journal of physiology Heart and circulatory physiology*. 2012;303 3:H315-22.
12. Dokainish H. Left ventricular diastolic function and dysfunction: Central role of echocardiography. *Glob Cardiol Sci Pract*. 2015;2015:3.



13. Carvalho J-C, Farand P, Do H, Brochu M-C, Bonenfant F, Lepage S. Effect of age and sex on echocardiographic left ventricular diastolic function parameters in patients with preserved ejection fraction and normal valvular function. *Cardiology Journal*. 2013;20(5):513-8.
14. Keller KM, Howlett SE. Sex Differences in the Biology and Pathology of the Aging Heart. *Canadian Journal of Cardiology*. 2016;32(9):1065-73.
15. Grassow L, Gröschel J, Saad H, Krüger LD, Kuhnt J, Müller M, et al. Sex-specific structural and functional cardiac remodeling during healthy aging assessed by cardiovascular magnetic resonance. *Clinical Research in Cardiology*. 2024.
16. Fleg JL, Strait J. Age-associated changes in cardiovascular structure and function: a fertile milieu for future disease. *Heart Fail Rev*. 2012;17(4-5):545-54.
17. Kwak HB. Aging, exercise, and extracellular matrix in the heart. *J Exerc Rehabil*. 2013;9(3):338-47.
18. Lieber SC, Aubry N, Pain J, Diaz G, Kim S-J, Vatner SF. Aging increases stiffness of cardiac myocytes measured by atomic force microscopy nanoindentation. *American Journal of Physiology-Heart and Circulatory Physiology*. 2004;287(2):H645-H51.
19. Tower J. Programmed cell death in aging. *Ageing Res Rev*. 2015;23(Pt A):90-100.
20. Biernacka A, Frangogiannis NG. Aging and cardiac fibrosis. *Aging and disease*. 2011;2(2):158.
21. Sanders D, Dudley M, Groban L. Diastolic dysfunction, cardiovascular aging, and the anesthesiologist. *Anesthesiol Clin*. 2009;27(3):497-517.
22. Parikh JD, Hollingsworth KG, Wallace D, Blamire AM, MacGowan GA. Normal age-related changes in left ventricular function: Role of afterload and subendocardial dysfunction. *Int J Cardiol*. 2016;223:306-12.
23. Bamaiyi AJN, Gavin R.*; Peterson, Vernice; Libhaber, Carlos D.; Sareli, Pinhas; Woodiwiss, Angela J. Limited contribution of left ventricular mass and remodelling to the impact of blood pressure on diastolic function in a community sample. *Journal of Hypertension*. 2019; 37(6):1191-9,.
24. de Simone G, Palmieri V. Diastolic dysfunction in arterial hypertension. *Journal of clinical hypertension (Greenwich, Conn)*. 2001;3(1):22-7.
25. Tran AH, Flynn JT, Becker RC, Daniels SR, Falkner BE, Ferguson M, et al. Subclinical systolic and diastolic dysfunction is evident in youth with elevated blood pressure. *Hypertension*. 2020;75(6):1551-6.
26. Fici F, Ural D, Tayfun S, Kozdag G, Facchetti R, Brambilla G, et al. Left ventricular diastolic dysfunction in newly diagnosed untreated hypertensive patients. *Blood pressure*. 2012;21(6):331-7.
27. Shim CY, Park S, Choi D, Yang W-I, Cho I-J, Choi E-Y, et al. Sex Differences in Central Hemodynamics and Their Relationship to Left Ventricular Diastolic Function. *Journal of the American College of Cardiology*. 2011;57(10):1226-33.